

Clinical

Motion sickness desensitisation:
a review of ADF experience of 'success'

Adrian Smith, Sonu Qurashi

Adrian Smith is a Specialist in Aerospace Medicine and research medical officer with the RAAF Institute of Aviation Medicine and HQ Forces Command.

Sonu Qurashi is a 4th year medical student at University of Adelaide who completed this project under the supervision of Dr Smith.

Motion sickness is a well-known phenomenon in aviation, predominantly amongst student pilots early in their flying training. As with many other air forces, the Royal Australian Air Force runs a motion sickness desensitization programme, comprising a 3-week course of repeated cross-coupled coriolis stimulation followed by a series of in-flight desensitisation manoeuvres. Progression through the cross-coupled stimulation phase with increasing rpm is determined by subjective comfort and vertigo time. Similar programmes around the world have reported 'success' rates between 70-90%; the AVMED desensitization programme claims a success rate of 91%. However, in reviewing different MSD programmes, the lack of an accepted definition of 'success' made it difficult to compare one programme with another. 'Success' can be defined as: completing the MSD protocol, returning to flying training, successfully completing basic flying training, or successful operational conversion onto a front-line aircraft. During the period 1997-2011, AVMED had performed 33 MSD courses on 30 aircrew. Of the 33 sessions, only 16 (48%) fully met the criteria for successful completion of the protocol, with another 13 (39%) being considered partial success and encouraged to return to flying, giving a notional 'success' rate between 48% and 87%. However, of the 20 student pilots followed up by interview, only 10 (50%) completed basic flying training, although the extent to which their failure was attributed to motion sickness could not be determined. Of the pilots who completed their flying training, 75% describe persistent operationally-significant motion sickness. This presentation will explore some of the reasons identified that can explain why so many people who passed the MSD protocol were unable to complete flying training, including interruption to flying training and infrequent provocative flying. Finally, this presentation will discuss an improvement to the way MSD candidates are followed up and managed when they return to flying, and improvements in the communication

between the flying instructors and AVMED in determining the extent to which motion sickness persists in student pilots despite them having undergone MSD, and the extent to which this may contribute to failure in flying training.

*Corresponding author: Adrian Smith
Email: adrian@pegasusaeromed.com*

Transfusion practice in critical care burns patients
– lessons for the military

CMDR Anthony Holley, Mr Nick Mckeough RN, LTCOL Michael Reade

Anthony is a dual qualified intensivist and emergency physician working at Royal Brisbane and Women's Hospital as a senior staff specialist. He is a senior lecturer with the University of Queensland Medical School and Chairman of Australia New Zealand Intensive Care Society, Queensland. Anthony was the recipient of the Matthew Davey Award that facilitated study at the Netherlands Military Blood Bank. He serves as a representative for the National Blood Authority Critical Care Expert Group. Anthony has deployed on a number of occasions including to Angola, Bougainville, East Timor, The South Pacific, The Persian Gulf and most recently to Afghanistan.

Introduction: There are no guidelines specifically directed to inform transfusion strategies in serious burn injury patients. This study was designed to describe the transfusion practice and evaluate the effects of transfusion on seriously burned patients. The study setting was a large university tertiary hospital intensive care unit, which is the only referral centre for burns injury in the state.

Methods: All burns patients admitted to a tertiary hospital referral centre intensive care over a 40 month period (9 December 2008 to 30 April 2012) were retrospectively identified. The patients were stratified on the bases of body surface area burned. The total number of blood product transfusions provided throughout the patient's admission was determined, including the number of units of blood transfused in and out of the operating theatre. Transfusion triggers were identified as active burn bleeding, active bleeding from "non burn" source, absolute haemoglobin level, and prophylactic transfusion prior to an invasive procedure or correction of coagulopathy (as diagnosed by International Normalised Ratio > 1.5 and Activated Partial Thromboplastin Time > 60 seconds). Outcome measurements included mortality, number of infections (diagnosed by the commencement of

antibiotics), and length of stay, duration of mechanical ventilation, acute kidney injury and number of surgical procedures.

Results: 158 burns patients were identified during the study period with a mean age of 41.6 years. The distribution according to burn surface area (BSA%) included 26 (0-9%), 21 (10-19%), 34 (20-29%), 23 (30-39%), 14 (40-49%), 10 (50-59%), 5 (60-69%), 1 (70%-79%), 10 (80-89%), 4 (90-100%) with 10 patients admitted with isolated airway burns. The mean APACHE II score was 17 +/- 7.5 with an average length of ICU stay of 10.99 days and a median length of stay of 5.95 days. The ICU mortality was 8.22%, but adjusted for "an intention to cure" was 0.63%. The mean number of transfusion episodes per patient for the duration of their ICU admission was 4.8 episodes (CI 1.66 95%). Most transfusions were performed in the intensive care unit as opposed to the operating theatre (3:1). The most frequent indication for transfusion were absolute haemoglobin correction, with prophylactic transfusion prior to an invasive procedure the next most common indication. Active bleeding from burn and "non burn" sites also constituted important indications. The mean INR triggering fresh frozen plasma transfusion was 1.6 +/- 0.92 (95%). The mean haemoglobin transfusion threshold was 89.1 +/- 25 g/dl and the mean platelet transfusion trigger was 66 x 10⁹/L +/- 63.

Conclusion: The most common indication for transfusion was anaemia. Practice was however highly variable. There also exists an association between units transfused and the number of infectious episodes, duration of stay, and duration of mechanical ventilation in patients with major burns, even after adjusting for burn severity. We were unable to demonstrate an association between the number of transfusions and mortality. Guidelines need to be established to inform rationale transfusion strategies for all blood product transfusions in the burn injury population.

*Corresponding author: Anthony Holley
Email: anthony_holley@health.qld.gov.au*

Lumbar multifidus muscle size is associated with lower limb overuse injuries in military specialist trainees

Mr Geoffrey Crowley, Dr Mark Creaby, Ms Sharne Neill, Prof Julie Hides, Dr Melinda Smith

Geoff is Senior Physiotherapist at the Amberley Health Centre, RAAF Base Amberley, Queensland.

Introduction: Lower limb overuse injuries are a significant cause of lost training time in military

recruits. Recent studies in other active populations provide evidence that the lumbopelvic and hip muscles may be associated with lower limb injuries. For example, decreased size of the lumbar multifidus muscle was predictive of lower limb injuries in football players and deficits in control of the trunk have been shown to be predictive of knee injuries in athletes. Similarly, a deficit in hip muscle strength has been associated with occurrence of lower limb injuries in college athletes. An important function of the lumbopelvic and hip muscles is effective load transfer and distribution of forces along the kinetic chain. Therefore the lumbopelvic and hip muscles play an important role in weight bearing activities. Given the relationship between lumbopelvic muscles, hip muscles and lower limb injuries reported in other active populations, the aim of this study was to compare lumbopelvic muscle size and hip muscle strength in military specialist trainees with and without a history of lower limb overuse injury.

Methods: Twenty-two male RAAF recruits participated in this study. All recruits successfully completed the 16-week RAAF basic training course. Injury history during the basic training course was established from military records. Participants were classified as either having sustained a lower limb overuse injury during basic training (LLOI), or not sustaining any injury during basic training (CTRL). Participants that sustained upper limb, trunk or acute injuries were excluded from this study. Bilateral ultrasound images of lumbar multifidus & quadratus lumborum were captured and measured to determine muscle cross sectional area (CSA, cm²). Maximal isometric strength tests of hip abduction and external rotation were also performed. Strength was reported as rotational force per kilogram of bodyweight (Nm.Kg). Participants were uninjured at the time of testing and all measures were taken prior to the commencement of Air-Ground-Defence specialist training. Independent t-tests were used to compare differences between the previously injured leg in the LLOI group and right leg in the CTRL group; an Alpha level of 0.05 was set.

Results: Age, height and weight were similar in the LLOI and CTRL groups (p>0.05). The CSA of lumbar multifidus was significantly greater in the CTRL group (9.41 ± 1.39 cm²) compared with the LLOI group (7.40 ± 0.48 cm²), p = 0.01. Conversely, there was no difference observed in the CSA of quadratus lumborum between groups (CTRL=7.23 ± 1.47 cm²; LLOI=6.30 ± 1.22 cm²), p = 0.26. No differences in hip abduction and hip external rotation strength were observed between the groups (p = 0.69 and 0.93 respectively).

Discussion: This study provides preliminary evidence that the CSA of the lumbar multifidus muscle is associated with lower limb overuse injuries during basic training. Given that decreased CSA of lumbar multifidus has been implicated in injury development in other populations, it would be prudent to screen for small lumbar multifidus CSA in injured recruits and consider targeted treatment of this muscle where appropriate. Future studies could assess whether an intervention program aimed at increasing the size of the lumbar multifidus muscle would reduce the likelihood of developing a lower limb overuse injury in Defence members.

Corresponding author: Geoffrey Crowley
Email: Gcrowls@hotmail.com

Penetrating Intracranial Injury caused by mud

Dr Martin Christie, M.B.Ch.B., Dip. Obs., F.R.A.C.S.

Martin trained in medicine initially in Rhodesia, and served as a Captain in the Rhodesian Army Medical Corps during his National Service during the early phases of enemy incursions. He then moved to Papua New Guinea, where he worked as a General Medical Officer for two years. This was followed firstly, by training in General Surgery and subsequently Neurosurgery, both in Dunedin, New Zealand. His later career included periods in the Republic of Kiribati, in Cambridge, UK, and Auckland, NZ. This was succeeded by eight years in Saudi Arabia, spanning both Gulf Wars, with exposure to casualties, as well as providing neurosurgical care for patients from the Bosnian conflict, and patients transferred to Saudi Arabia from the civil war in Yemen. He moved to Coventry, UK, working as a consultant neurosurgeon in the NHS. He has served four tours of duty in Afghanistan as neurosurgeon to the NATO coalition, after retiring from the NHS. More recently he has instructed military doctors in the new state of Southern Sudan. Recently retired, he lives in Sydney.

The proliferation of explosive devices cached beneath the earth leads to predictable patterns of injury. In the explosion multiple metallic fragments are sprayed out, together with the overlying soil and its various components. Among the consequences for persons near to such detonations is the penetration of the body by these soil components as a result of the kinetic energy imparted to them from the explosion. The velocity and mass of the individual particles have a direct bearing on the damage sustained in human tissue during the abrupt process of stopping so that a piece of gravel travelling at speed will do more harm than a grain of sand.

A case was recently encountered during the current conflict in Afghanistan, where it appears that the explosion resulting from an improvised explosive device (IED) triggered by a pressure plate, caused a clod of overlying mud to possess sufficient penetrating power to fracture the skull and enter the cranial cavity. The emergency surgical treatment and difficulties in recognition of this injury are described. The rarity of this mechanism of injury is in contrast to the patterns of injury usually sustained in such incidents. Expeditious intervention is crucial in dealing with this type of contamination in the intracranial extradural space and serves to emphasise the importance of placing neurosurgeons amongst the personnel closest to the battlefield.

Corresponding author: Martin Christie
Email: barmarbaratin@hotmail.com

Tendinopathies, Enthesopathies, Overuse Injuries and Alphabet Soup. Where is the Science?

Tony Delaney

Dr Delaney conducts a specialist sports medicine practice at Narrabeen Sports Medicine Centre, Sydney Academy of Sport, Narrabeen.

He was the visiting senior specialist in sports and rehabilitation medicine to the Fleet Base East Medical Centre, HMAS Kuttabul and IHSB, Holsworthy Military Area.

He was principal of the Medical Centre, Charlotte Pass, Snowy Mountains 1974-92 winter seasons. The practice is snowbound and provided acute trauma, radiology, emergency, general and sports medicine service Charlotte Pass Ski Patrol coordinated with Snowy Mountain Search and Rescue services.

His Special Interests are:

- *Biomechanics, overuse and acute injuries of the spine, lower limb and upper limb.*
- *Human performance, physiology and medicine in the environments of heat, cold, high altitude and underwater.*

Military Medicine.

How do we decipher Corticosteroid injections, Plasma rich protein, Autologous blood injection. Nitric Oxide patches, Low Intensity Pulsed Ultrasound, Extracorporeal Shock Wave Therapy, Faradics, Interferential, Acupuncture, Dry needling for treatment of chronic and acute overuse injuries. The scary thing is that most modalities are a waste of time and money. The clinician who addresses the biomechanical causes of these syndromes will

achieve optimum results. This presentation will provide a summary of current best practice for tennis elbow, shoulder impingement, patellar and Achilles

tendinosis, plantar fasciitis. All of the things that frustrate you in clinical practice in Military Medicine.

Corresponding author: Tony Delaney

Email: drtonydelaney@hotmail.com

Mental Health

Current Themes in Operational Mental Health

Maureen Montalban

MAJ Maureen Montalban is currently the Officer Commanding of the 1st Psychology Unit, Sydney Detachment. She joined the Regular Army in 2004 as a psychology officer and throughout her career has worked in research, assessment, counselling, training and operational psychology. MAJ Montalban has deployed in support of ADF personnel to Operation CATALYST, Operation ASTUTE, Operation ANODE and Operation SLIPPER. She is currently in her final year of her Master of Psychology (Health).

This presentation will cover the key operational mental health themes from all areas of operation through discussion of both the questionnaire data obtained through the Return to Australia Psychological Screens (RtAPS) and Special Screens; as well as through referrals on operations, command liaison and early intervention activities. These screening tools and interventions not only assist with the mental health surveillance of the workforce, they also provide a mechanism for identifying groups who may be at high risk of exposure to stress or trauma. The ongoing work with groups identified as being "at risk", including specific interventions to assist with mental well-being, will be discussed.

Corresponding author: Maureen Montalban

Email: maureen.montalban@defence.gov.au

Enhancing capability and resilience through Positive Psychology

Mark Mathieson (Major – Army Reserve)

Mark is a registered Psychologist and currently serving Major in the Army Reserve – Australian Army Psychology Corps.

Mark has experienced a broad variety of roles within the industrial psychology sphere and has served the ADF for over 15 years including six years supporting Special Forces and numerous deployments. Mark has designed, managed and successfully achieved outcomes on diverse and complex civilian projects

including green-field LNG sites in Papua New Guinea, mining companies across Australia and in Laos and Timor Leste, utility companies, NGO's and with individuals and organisations working in the sustainability sector.

The last decade has seen a rapid growth in the field of Positive Psychology, including the application of Positive Psychology frameworks and tools in the areas of education, employment and individual psychotherapy. Outcomes in these areas have shown a remarkable capacity to build both excellence in performance, but also robustness and resilience in individuals who are provided with the tools and knowledge to buffer against the demands of difficult times. Many organisations are now turning to proactive models of intervention in order to 'front load' individuals and teams with resilience and a capacity to cope. Whilst the natural assumption may be that building resilience is simply the removal or reduction of existing stressors or an enhancement of an individual's capacity to move themselves from negative functioning back to 'normal functioning' when times are bad, Positive Psychology theory and research suggests that a different set of variables need to be developed if individuals are to move towards 'maximal functioning'. In this interesting and though provoking presentation, Mark will outline the basis and theoretical models underpinning Positive Psychology and how they can be applied to build resilience and coping capacity in individuals within the ADF and Veterans communities. Based on his 15 years of experience in working with individuals and organisations selecting, deploying and returning people from isolated and remote environments including the ADF in general, on operations, Special Forces Selection Courses and the Australian National Antarctic Research Expedition, Mark will use real life examples of how Positive Psychology can significantly enhance capacity, protect against negative mental health outcomes and ultimately – enhance capability.

Corresponding author: Mark Mathieson

Email: mmathieson.psychedup@gmail.com

Pre-deployment BattleSMART: The development and evolution of psychological resilience training for ADF personnel preparing for operations

Nicole Sadler

COL Nicole Sadler is currently the Director Strategic and Operational Mental Health within Joint Health Command. COL Sadler joined the Regular Army in 1994 as a psychology officer and throughout her career has worked in recruitment, assessment, counselling, training, policy development and operational psychology. She was the Commanding Officer of 1st Psychology Unit Jan 10-Aug 12. COL Sadler has deployed in support of ADF personnel to Operation BEL ISI, Operation SUMATRA ASSIST, Operation CATALYST, Operation ASTUTE and Operation SLIPPER. She completed the Australian Command and Staff Course in 2004 and was awarded a Master of Psychology (Clinical) degree in 2005.

Following the Dunt Review and the Government's 2007 commitment to a Mental Health Lifecycle Package for ADF personnel, the ADF commenced rolling out a comprehensive whole of career resilience training program named BattleSMART (Self Management and Resilience Training) in 2009. BattleSMART is a modularised educational program that operates across the ADF, teaching resilience training at key points throughout a member's career. It is a preventive program designed to build psychological resilience by teaching ADF personnel to test and adjust their physical, cognitive, emotional and behavioural reactions to stressful situations as required, to optimise their performance.

The pre-deployment BattleSMART component was developed by staff from the Directorate of Strategic and Operational Mental Health, Joint Health Command and 1st Psychology Unit and was first trialled in 2010. Pre-deployment BattleSMART focuses on the specific challenges likely to be faced leading up to and during deployment, and promotes both individual and collective optimal performance through the enhancement of individual coping. Training delivery is a combination of instructor led theory presentation and practical exercises, and the results of recent evaluations conducted with soldiers preparing for Operation SLIPPER have been very positive. This presentation will outline the development and evolution of pre-deployment BattleSMART, the program content, evaluation outcomes, and the mechanisms for continuously reviewing and updating the content to ensure relevant information is presented to ADF personnel as they prepare for the different Areas of Operations.

*Corresponding author: Nicole Sadler
Email: nicole.sadler@defence.gov.au*

RESET – Recognising Early Signs of Emerging Trauma: An indicated prevention program for PTSD

Dr Stephen Rayner & Ms Jane Nursey

Stephen Rayner is contracted to Joint Health Command as Programs Manager at the ADF Centre for Mental Health in Sydney. He is a clinical psychologist with over 20 years experience working for Defence as a public servant, contractor and Naval Officer and has published several papers and a book chapters on military mental health.

Jane Nursey is a Senior Clinical Specialist at the Australian Centre for Posttraumatic Mental Health (ACPMH). She is a Clinical Neuropsychologist with over 20 years experience working in public and private health and mental health services. This includes many years experience in delivering PTSD treatment programs for Veterans, current serving members, emergency services personnel and members of the community. Her work at ACPMH involves the translation of research findings into evidence based policy and service development advice as well as the training of clinicians in evidence based practice for the treatment of trauma related mental health disorders.

The ADF has recognised a gap in the provision of a comprehensive program of interventions for Posttraumatic Stress Disorder (PTSD); and the ADF recognises that ADF personnel are at increased risk of developing PTSD due to the nature of military service. While there are recognised general and targeted prevention strategies, and treatment for diagnosed disorders, there exists a gap for indicated prevention strategies – those targeting people who have early and detectable indicators foreshadowing emerging PTSD, but who do not yet have a diagnosable disorder. Indicated prevention strategies have been applied to depression and anxiety, but there is limited acceptance of this type of intervention for PTSD. The ADF, in collaboration with the Australian Centre for Posttraumatic Mental Health has developed an intervention program to intervene with ADF members with emerging, but not yet diagnosable, PTSD. The program aims to increase resilience to PTSD, to manage existing and potential symptoms of PTSD, reduce the likelihood of progression to disorder, and improve quality of life. The program is evidence-informed and based on existing practice guidelines. The program has been piloted recently, and this paper will describe the development of the program and report on preliminary evaluation data from the pilot program.

*Corresponding author: Stephen Rayner
Email: stephen.rayner4@defence.gov.au*

Mild Traumatic Brain Injury (MTBI) in the Australian Defence Force: Results from the 2010 ADF Mental Health Prevalence and Wellbeing Dataset

Alexander McFarlane

Professor Alexander McFarlane currently the Head of the University of Adelaide Centre for Traumatic Stress Studies and is chief investigator on the Mental Health Prevalence Study. He qualified in Medicine in 1976 with Honours and completed his specialist training in Psychiatry in 1980. In 1990 he was awarded the degree of Doctor of Medicine based on his longitudinal research into the aftermath of the Ash Wednesday Bushfires in South Australia. Subsequently he became an international expert in the field of the impact of disasters and posttraumatic stress disorder (PTSD). He has published over 250 articles and chapters in various refereed journals and has co-edited three books. He currently holds the rank of Group Captain in the RAAF specialist reserve. In 2011 he received the Officer of the Order of Australia award in the Australia Day Honours List. The award recognises outstanding contributions to medical research in the field of psychiatry, particularly posttraumatic stress disorders, to veterans' mental health management, and as an author.

The occurrence of mild traumatic brain injury (mTBI) has attracted much attention in the research

literature, as well as public domains (such as the media), in recent years. This has emerged due to a suspected increase in mTBI, thought to be the result of increased use of explosive devices in combat in the last decade. Accurate prevalence estimates of MTBI are extremely difficult to ascertain given the non-specific nature of post-concussive symptoms, measurement issues (to determine the degree of combat exposure), variation in deployment length as well as cultural differences such as compensation practices and healthcare systems across countries. Understanding mTBI in the context of deployment is important owing to the implications for healthcare provision, deployability status and compensation for affected veterans. As such, there is a distinct lack of epidemiological estimates of mTBI in military populations, including the Australian Defence Force that needs to be addressed. This presentation will examine the lifetime prevalence of self-reported head injury in a representative sample of the ADF. Mechanisms of injury, frequency of reported post-injury symptoms, differences between deployed and non-deployed groups, and relationships between injury and various psychiatric disorders will also be explored.

Corresponding author: Alexander McFarlane

Email: alexander.mcfarlane@adelaide.edu.au

Clinical Training

Development of a medical officer Pharmacology Training Package for the RAAF – Update

SQNLDR Michael Lumsden-Steel

SQNLDR Michael Lumsden-Steel is an EX APF RAAF Medical Officer and now RAAF SR whilst undertaking anaesthetic training on the Tasmanian Anaesthetic Training Rotational Scheme.

Dr Michael Lumsden-Steel will present an update on the development a medical officer pharmacology training package for the Air Force.

Corresponding author: Michael Lumsden-Steel

Email: mc.ls@me.com

Maintaining connectedness and skills in the geographically dispersed clinical workforce

James Ross, Meredith Leonard

Dr James Ross is the Medical Director of Aspen Medical, based in Canberra. He is a Fellow of Occupational and environmental Medicine, Public Health Medicine and Aerospace Medicine, and the

current President of the Australasian Faculty of Occupational and Environmental Medicine of the RACP. He is the foundation president of AMMA 1991-1995. He had 26 years in the permanent Air force, and is currently Group Captain in the RAAF specialist reserve.

As a major private health service provider, with clinical staff located across all Australian states and territories and internationally, Aspen Medical is committed to providing its geographically dispersed clinical workforce with the support and resources to provide consistently high quality health care services. Operating in remote locations does provide challenges to ensuring the health professionals maintain a connection with their employer, and their processes, and of maintaining clinical currency within the workforce.

Aspen Medical has developed and implemented a web based solution to address such challenges. Through this platform staff can access company policies, resources, and training opportunities. This solution has been well received by staff. However,

such a system does raise its own challenges from which lessons must be learnt in order to progress forwards. User accessibility and technical ability have proven to be the greatest hurdles.

The prioritisation of training resource development can become overwhelming. In the first instance, Aspen Medical has chosen to focus its training development energies around priority areas as identified by the Australian Commission on Quality and Safety in Healthcare. Recognised knowledge deficits or skill gaps and demand from customers have also been considered in developing a company training development plan. Examples are Clinical Practice Guideline understanding and assessment; drug calculations, common medication errors and hand hygiene.

One of the great successes of this system has been the centralisation of training records and the ability to ensure the standardised delivery of training across all sectors of our geographically dispersed workforce. This is also essential in permitting compliance with Registered Training Organisation certification, and with obtaining CPD points from organisations such as RACGP.

This presentation will demonstrate the current capabilities of the system, planned improvements and lessons from Aspen's experience in establishing such a system

Corresponding author: James Ross
Email: james@rahc.com.au

Northern Territory ADF Clinical Placement Program

Jessica Burton and Bronte Douglas

Jessica Burton spent 8 years in the RAAF as a Nursing Officer posted to 1ATHS in Amberley and 3EHS Detachment Darwin. She transferred to the Active Reserves in 2010 to be able to clinically up skill in the Intensive Care environment and has since completed her Graduate Certificate in Critical Care Nursing. She was employed by the National Critical Care and Trauma Response Centre, delegated by the NT Department of Health, in 2011 to create and organise the NT ADF Clinical Placement Program.

The Northern Territory has traditionally had a good relationship with the Australian Defence Force (ADF) and this is fostered in many forms. Since 1999 the Northern Territory Department of Health (NT DoH) and ADF has had a Memorandum of Agreement (MOA) in place, which includes the ability to place ADF Health Personnel in a NT DoH facility on clinical

placement. In 2011 a variation to the MOA was signed between the two parties and in May 2011 the NT DoH created the ADF Clinical Facilitator position; to be managed daily by the National Critical Care and Trauma Response Centre (NCCTRC). The position was created as a Nursing Level 4 (Nursing Educator Level) at 0.4FTE; the requirement was to be filled by a NT DoH member who had working knowledge of both NT DoH and ADF Health. This ensured the person was able to understand the requirements of and appropriately converse with NT DoH Managers and ADF Managers.

Since the creation of the ADF Clinical Facilitator position the responsibilities have included:

- Creating, organising and facilitating Clinical Placements for all ADF Health Personnel within NT DoH facilities.

Secondary Duties include:

- Fostering and maintaining good working relationships with NT DoH clinical areas.
- Seeking and creating opportunities for ADF Health personnel in clinical areas, education and other forms of training.
- Become a point of contact regarding all things ADF within the NT DoH.

Creation of the Clinical Placements

The NT ADF Clinical Placement Program was initially modelled off the Princess Alexandra Hospital rotations to Royal Darwin Hospital. The model deems ADF members supernumerary to the clinical areas numbers, however take a (patient) load once orientated and confident. Since the commencement of the program we have had Medical Officers, Nursing Officers, Medical Technicians and a Pharmacist conduct Clinical Placements. The areas have included Operating Theatre (including Anaesthetics), Emergency, Intensive Care/High Dependency, Cardiac Care, Acute Surgical (including Burns), General Surgical and Acute/General Medical.

The following Lessons have been learnt since the commencement of the placements:

- The ADF need an appropriate Clinical Uniform for placements
- Appropriate recalling of personnel
- Appropriate lengths of placements
- Infection Control – feeding information back into the ADF Health system
- Clinical area placement caveats – including Endorsed Enrolled Nurses

Future Ideas

- Planning a Predeployment Clinical Placement Stream – The clientele issues, clinical cases and language barriers are similar to what ADF personnel will experience on deployment or humanitarian missions.
- Clinical Placements for interstate ADF Health Personnel
- Provide Postgraduate Training for ADF Health Personnel

Overall, the ADF Clinical Placement Program has been very well received within NT DoH facilities.

Corresponding author: Jessica Burton

Email: jessica.burton@nt.gov.au

Reinvigorating CBRN Health Training for Defence Personnel- a new perspective

Kylie Douglas

Kylie Douglas is currently the Training Development officer for the Directorate of Workforce Development and Training at Joint Health Command. She is involved in a number of Joint Health training initiatives including the CBRN Health Course, Military Anaesthetics Course and the training requirements for the new Garrison Health Services contract. In addition Kylie is the Defence liaison officer for the Community Services and Health Industry Skills Council.

Ms. Douglas has been working in Defence for the last fifteen years and previously served as a RAAF Officer. In that capacity she was involved in the establishment of the ADF Medics training continuum, the ongoing AME training requirements for ADF health personnel and the establishment of the Military Critical Care AME Team (MCAT) course. She has both clinical and operational health experience and holds a Diploma of Training and Assessment Systems in addition to a Masters of International Public Health.

Chemical, Biological, Radiological and Nuclear (CBRN) health concerns from an ADF perspective have evolved considerably since the proliferation of chemical defence weapons in WWII. Whilst the modern day threat of CBRN weaponry is arguably low, the need for Defence health personnel to be prepared for casualties with CBRN related injuries is steadily climbing. Whole Of Government responses to global crises sees our health personnel involved in numerous CBRN type incidents, such as Fukushima, whilst at home the clean up from WWII chemical munition dumps continues. How we prepare our health professionals to work in challenging

conditions, recognize broad symptomatology, provide advice to commanders on public health concerns and work effectively with multiple stakeholders is paramount to a successful mission.

In the last five years competing Defence priorities has resulted in the demise of corporate knowledge surrounding CBRN health related medicine and broader CBRN awareness across Defence. The Services in recognizing this shortfall are reinvigorating CBRN capability with Joint Health Command (JHC), Special Operations Engineer Regiment (SOER) and the Army Logistic Training Centre (ALTC) re-establishing CBRN health specific training for our personnel.

This presentation will examine the broader Defence initiatives in relation to CBRN capability and explore the challenges for health personnel in meeting both individual and collective needs. To do this, the training and outcomes of the CBRN Health course pilot conducted at SOER in September will be discussed along with the current direction of CBRN health globally.

Corresponding author: Kylie Douglas

Email: kylie.douglas3@defence.gov.au

Care in Combat Training for Defence

Glenn Keys

Glenn Keys is the CEO and Managing Director of Aspen Medical. Glenn's career covers a broad range of businesses, from start-up's to US multi-nationals. After a distinguished career in the Australian Army, where he covered a range of tasks, from test flying to engineering and logistics support for Army aircraft, Glenn was responsible for the establishment of a number of new businesses, either as start-ups or as new business units in global corporations. Glenn has led Aspen Medical from its founding just over nine years ago to today when Aspen Medical is a substantial international business with a presence in the Asia, the Pacific, USA, Canada, Papua New Guinea, Timor Leste, the Middle East, Australia and the United Kingdom. Glenn also has a strong sense of community involvement, working closely with organisations such as the ACT Down Syndrome Association, BLITS and the Special Olympics ACT Region.

Outlines the background for the course, and development of key curriculum points, including decision to include tourniquet as a method of increasing survival at the sacrifice of the limb. Outlines staff selection, including decisions to use only staff who have seen service under fire in areas

of deployment. Covers equipment selection, and notes requirement for equipment that can meet the demands of the training environment. Covers expansion of the course to include PMV's and environmental effects to increase realism of course. Finally outlines areas of continual development of

the course to increase effectiveness in a changing combat environment, including eliciting feedback on return from deployments.

Corresponding author: Glenn Keys

Email: gkeys@aspenmedical.com.au

Veteran's Health

Considerations in the dental treatment for Vietnam war veterans with PTSD

Adj. A/Prof Norton Duckmanton, OAM

Prof. Duckmanton is a Prosthodontist at Sydney Dental Hospital treating mostly DVA patients with implant supported prostheses. He is also teaching in the Prosthodontics specialty programme within Sydney University. Upon graduation he conducted practice as a dentist for 10 years when he entered the Faculty of Dentistry, retiring 25 years later. He served during that time as a visiting Professor in Prosthodontics at North Western University Dental School in Chicago on two occasions. He was flying as a Navigator in WWII in the SWPA. He was a member of the RAAF Reserve from 1947 to 1982 retiring as a Consultant in Prosthodontics with the rank of Group Captain.

Dental treatment for Vietnam Veterans is complicated by many factors; some of which are parafunctional muscle activity and nocturnal bruxing; which leads to fracture of restorations and tooth structure and masticatory muscle spasm, pain and dysfunction.

A reduction in salivary flow, as a side effect from the prescription of anti depressant medication predisposes accelerated recurrent marginal caries attack, and a reduction in the pain tolerance of the soft tissues. Alcohol and drug abuse, expressions of anger, low self esteem and breakdown of family relationships often leads to a poor compliance in home care instructions.

A multidisciplinary team composing the patient's physician, psychiatrist, clinical psychologist, social worker and dentist is considered to be an ideal composition.

An aggressive oral hygiene and caries preventive program should be initiated at the outset, as a protection from new

And recurrent caries and from periodontal health breakdown. The use of specially designed trays for the administration of remineralising and cariostatic medicaments is also desirable.

An initial soft mouth guard/splint to limit the effect of bruxism while treatment is in progress is recommended.

Restorations and prosthodontics appliances should be of a robust and simple design in order to withstand the loads generated by nocturnal and diurnal muscle parafunction and bruxism. These appliances are also simple to maintain and repair.

The use of multiple full coverage crowns to prevent tooth and restoration fracture may be subject to marginal recurrent caries where compliance of a strict oral hygiene is low, and in such cases should be prescribed with caution.

The provision of a protective occlusal splint is necessary at the completion of treatment, to protect the dentition and restorations from the effects of parafunctional activity and bruxism. Where there is wide spread periodontal bone loss, active caries and low compliance of oral hygiene instructions, it can be predicted that tooth loss will be accelerated as time progresses. It is suggested that a regular maxillary complete denture opposing a mandibular complete overdenture, supported and retained by two implants, would be an appropriate final restoration, which has proved to be an effective means of providing an oral restoration requiring only simple, regular maintenance.

Corresponding author: Norton Duckmanton

Email: norton.duckmanton@email.cs.nsw.gov.au

The health and wellbeing of Australia's female Vietnam and contemporary veterans

Samantha Crompvoets, PhD

Dr Samantha Crompvoets, BSC (hons) Melb PhD ANU, is a Sociologist and Research Fellow in the ANU Medical school. Dr Crompvoets is Chief Investigator on a number of DVA and Defence funded projects examining issues of gender, workforce, identity, culture, health and wellbeing. Her current projects include examining the health and wellbeing of female veterans, and the history, identity and wellbeing of ADF reservists. Other research in particular includes

an analysis of the free text responses in the recent ADF census and reserve attitude survey, compiling a history of the Defence Reserves Association and analysing the marketing and recruitment campaigns aimed at Reserves since Federation.

This paper presents the findings from a three year DVA funded study into the health and wellbeing of female veterans. Empirical data was collected through face to face interviews with 90 veterans and stakeholders in female veterans' health and care. Participants gave complex accounts of their experiences of deployment, broader military career and life since discharging from the ADF. This complexity has been summarised and translated into evidence for action to improve the health and wellbeing of all female veterans. This presentation will overview barriers to accessing existing support services for female veterans, gaps in available and appropriate information, resources and DVA policies for female veterans, and gaps in knowledge of female veterans that impact health and wellbeing and service provision.

*Corresponding author: Samantha Crompvoets
Email: samantha.crompvoets@anu.edu.au*

Cancer incidence and mortality in Australian Gulf War veterans

Dr Jillian Ikin, Ms Stella-May Gwini, Dr Helen Kelsall, Ms Breanna Wright, Prof Malcolm Sim

Dr Jillian Ikin is a Research Fellow at the Monash Centre for Occupational and Environmental Health, Department of Epidemiology & Preventive Medicine, Monash University. She was an investigator on the baseline Australian Gulf War Veterans' Health Study 2001-2003, the Korean War Veterans' Health Study 2005 and is again an investigator on the current Follow Up Health Study of the Gulf War veterans and their military comparison group. Dr Ikin completed her PhD on war stressors and mental health outcomes in the Australian Gulf War and Korean War study cohorts.

Background: A previous linkage study of Australian Gulf War Veterans (GWVs) found no statistically significant excess of cancer up to the end of 1998 or mortality up to the end of 2000, although the numbers of deaths and cancers were small. In 2012 we have repeated the cancer and mortality linkage for the same study cohort.

Aims: The analyses aimed to investigate whether veterans of the 1991 GW have an excess risk of death or of developing cancer compared with a military comparison group (CG) and compared with the Australian general population.

Methods: The study cohort of 1,871 GWVs and 2,923 CG members was linked to the National Death Index and Australian Cancer Database. The observation period was 1st January 1991 to 30th of November 2007 for mortality and 1st January 1991 to 31st of December 2008 for cancer incidence. Australian general population data was used to calculate the expected numbers of deaths or cancers in each year of follow up, and these were compared with the cohort observed numbers to calculate standardized incidence and mortality rates (SIR and SMRs). The rate of death or cancer in the GWVs was compared with that in the CG using rate ratios (RR) adjusted for service type, rank at the time of the GW deployment and age.

Results: There were 42 deaths in the GWV group and 51 in the CG. The all-cause SMR was 83 (95% CI 62-113) for the GWVs and 64 (95% CI 49-84) for the CG, with neither finding statistically significantly different to the general population. There was no statistically significant excess of all-cause mortality in GWVs compared with the CG, with a RR of 127 (95% CI 85-192). The SMR for external causes of deaths was 74 (95% CI 46-121) for GWVs and 63 (95% CI 41-97) for the CG, not statistically significantly different from the general population in either group. The GWVs had a greater risk of deaths from external causes (RR= 117 95% CI 61-224) although this was also not statistically significant. The SIR for all malignant neoplasms in the GWVs was 99 (95% CI 76-133) and for the CG the SIR was 83 (95% CI 65-107). When compared with the CG, the rate of all malignant neoplasms among the GWVs was not significantly different to that of the CG group (RR 116 95% CI 80-167).

Conclusion: As of the end of 2008 for cancer and the end of 2007 for deaths, the results for Australian GWVs are not statistically different from those in the general community or the military CG. There was insufficient numbers to permit analyses of specific causes of death or specific cancer types. However, the cohort is still young and the number of person years low, indicating the need to follow this cohort into the future.

*Corresponding author: Jillian Ikin
Email: jill.blackman@monash.edu*

Risk and resilience in Australian military families

Dr Annabel McGuire, Ms Catherine Runge, Dr Renee Anderson, Ms Katrina Bredhauer, Professor Annette Dobson, A/Prof Peter Nasveld

Dr Annabel McGuire joined the Centre for Military and Veterans' Health (CMVH) in January 2007 as a Research Fellow within the Deployment Health Surveillance Program. Currently, she is the Chief Investigator on the Timor-Leste Family Study. During her time with CMVH, Annabel has been co-recipient of four Defence and DVA grants, produced six peer-reviewed journal articles and more than 20 reports to Defence & DVA. Annabel has been co-author of 25 conference presentations, presenting nine of these.

Risk factors are conditions or variables associated with a lower likelihood of positive outcomes and a higher likelihood of negative or socially undesirable outcomes. Protective factors have the reverse effect: they enhance the likelihood of positive outcomes and lessen the likelihood of negative consequences from exposure to risk^[1]. The same factor may be either risk or protective. Resilience is about responding and adapting to crises and adversity, and recovering and growing from these experiences^[2]. Individual strengths, family strengths and community supports all play a role in resilient families^[3].

The Australian Department of Veterans' Affairs funded a large cross-sectional quantitative study as part of the larger Family Study Program. Responses were collected from Australian veterans and partners (N>4,000) using validated measures of physical (SF-12), mental (K10, SF-12, PCL-C), family (FACES-IV) and child (SDQ) health. Data on risk and protective

factors was also collected including measures on how people cope (BriefCOPE), the social support they access (DUSOCS), intimate partner violence (WAST) and their overall relationship satisfaction (QRI).

The presentation will investigate potential risk and protective factors associated with the physical, mental, and family health of partners and children including family functioning, coping, social support, quality of relationship, and intimate partner violence. Risk and protective factors may exacerbate or ameliorate effects associated with military life for partners and children. The findings support aspects of prior research particularly from the United States and provide new insights on the health and wellbeing of Australian military families and their family dynamics. Factors associated with risk and resilience for military families may be amenable to policy and practice interventions.

References:

1. Jessor, R., Turbin, M. Z., & Costa, F. M. (1998). Risk and protection in successful outcomes among disadvantaged adolescents. *Applied Developmental Science*, 2(4), 194-208
2. Walsh, F. (2003). Family Resilience: A Framework for Clinical Practice. *Family Process*, 42(1), 1-18. doi:10.1111/j.1545-5300.2003.00001.x
3. McCubbin, H. I., & McCubbin, M. A. (1988). Typologies of resilient families: Emerging roles of social class and ethnicity. *Family Relations*, 37(3), 247-254

Corresponding author: Annabel McGuire

Email: a.mcguire@uq.edu.au

Ethics and Legal Aspects

When is it permitted to break medical in confidence? a discussion of the ethical issues for defence medical staff in a case of codeine addiction

SQNLDR Michael Clements

SQNLDR Clements is the Health Centre Manager at Townsville Health Centre at RAAF Base Townsville. Since completing the Graduate Medical Scheme at University of Sydney with initial training at Royal Prince Alfred Hospital in Sydney and in Geraldton, WA, he was posted to RAAF Base Tindal and then to the United Kingdom for the Diploma in Aviation Medicine. After a short period as the Officer in Charge of the 1 EHS Detachment at Townsville he was posted to his current position in Joint Health Command. Current research interests include Occupational medicine and Medical administration.

Codeine addiction is a commonly occurring substance use disorder in the Australian general population that is facilitated by the easy access to over the counter preparations. Despite the increasing regulation in which pharmacists are allowed to dispense the medication, the medical literature is growing about the impact of the addictions with life threatening gastrointestinal disease, renal failure, anaemia and severe hypokalaemia. In this case study what had initially appeared to be a diagnostic conundrum of life threatening anaemia on a background of gastric erosions later revealed itself as a substance use disorder.

The diagnosis and rehabilitation plan required careful balancing of the needs of the individual and the needs of the Commonwealth. In this particular case, we found that medical advice and management

alone was insufficient to generate a change in behaviour, and the inclusion of the member's chain of command was crucial to the patient engaging in the therapeutic process. The way that the matter has been handled in a medical, administrative and disciplinary sense have revealed a complex but compassionate, positive and patient centered approach that reveals values in the best traditions of the ADF.

The frontline medical officer must maintain a high degree of suspicion of codeine addiction when managing patients with some of the non-specific comorbidities of this substance use disorder. Once suspected or revealed, the complex interaction that is required between the medical officer and the chain of command needs to be well thought out and planned in order to provide the best outcomes for the patient and the Commonwealth.

Corresponding author: Michael Clements

Email: michael.clements@defence.gov.au

Statements of Principles, process and challenges with translating epidemiology into legal instruments.

Justine Ward, Paul Murdoch, Nick Saunders

Justine Ward is a public health physician and Principal Medical Officer of the Repatriation Medical Authority Secretariat.

Since 1994 the system for establishing liability for compensation for most military service in Australia has been based on Statements of Principles (SoPs). SoPs are legal instruments which set out the factors that must exist to cause a particular kind of disease, injury or death that could be related to eligible service, based on sound medical-scientific evidence (SMSE). Decision-makers determining claims from individuals cannot accept a claim unless it meets at least one factor, and associated 'dose'. The SoPs are determined by the Repatriation Medical Authority (RMA). The RMA consists of five practitioners eminent in the field of medical science, supported by a secretariat. SoPs are determined by assessing the SMSE for evidence of a causal association between potential risk factors and a given disease. Up to the end of the 2011 financial year, the RMA had produced 1833 instruments covering 304 conditions. Conditions covered are wide ranging in nature and include chronic diseases, infections, psychiatric conditions, injuries, overuse syndromes and cancers.

The legislation requires that the condition meet the Veterans' Entitlements Act 1986 (VEA) definition of "disease" and that the RMA assess causation using

published peer-reviewed evidence and standard epidemiological criteria. The latter are based on the Bradford-Hill criteria, and include temporality, strength of association, observation of a dose-response effect, biological plausibility, consistency with other evidence, and absence of alternative explanations for an association (chance, bias or confounding). Consideration of the quality of the published evidence is an important part of the critical appraisal process. The assessment of the evidence is complicated by the fact that there are two standards of proof, reasonable hypothesis (RH) and balance of probabilities (BoP). For the RH standard, the SMSE has to indicate or point to a reasonable hypothesis of a causal association between the factor and disease. For the BoP standard, the SMSE has to show that it is more probable than not that the factor is causally related to the disease. While the same body of SMSE is used for each disease, the factors and related doses may vary between the 2 SOPs.

In experimental conditions factors other than the association of interest can be controlled, but epidemiology by its nature is an imperfect science, using data from free-ranging human beings. This paper will give examples of associations which are more difficult to assess for various reasons, including those which are of particular interest to veterans.

Problems with the evidence include lack of information concerning the association in question, poor exposure assessment, mismatch between the evidence and popular beliefs, reverse causation, recall bias, variable disease definitions and poor control for confounding. A wealth of new data on genetic risk factors is being generated, and the challenge of applying information on genetic risk factors to SoPs will also be discussed.

Corresponding author: Justine Ward

Email: justine.ward@rma.gov.au

Credentialing and the law

Ms Yolanda Kuruc

The process of credentialing health professionals is more than a tedious rubber-stamping process but rather an unparalleled opportunity to effect patient safety. Credentialing is described in the 2004 Australian Council on Safety and Quality Council (now the 'Commission') publication 'Standard for Credentialing and Defining the Scope of Clinical Practice as the 'formal process used to verify the qualifications, experience professional standing and other relevant professional attributes of medical practitioners for the purpose of forming a view about their competence, performance and professional

suitability to provide safe, high quality health care services within specific organisational environments’.

Rigorous and robust credentialing processes can protect patients from potentially avoidable harm. The enquires into the notorious appointments of Dr Graeme Reeves and Dr Jayant Patel provide valuable insight into what expectations patients may reasonably have of the credentialing processes of health care facilities. The failings by the responsible personnel to adequately assess the qualifications and experience of these two doctors, and the inadequate response to ‘red flags’ in the doctors’ applications for employment, is described in the Davies and 2008 ‘Garling Report’ into the circumstances surrounding the appointment of Dr Graeme Reeves. This paper exams the duties owed by health care facilities to patients in relation to credentialing the health professionals.

Corresponding author: Yolanda Kuruc

Email: ykuruc@aspenmedical.com.au

International Humanitarian Law / Law of Armed Conflict in relation to health personnel. Why do we still not understand?

David Thompson

LCDR David Thompson RANR, is a Reserve General Practitioner. He has recently transferred from Army Reserve to RANR. His operational service includes 2 deployments to Afghanistan: 1stly, as part of the forward rotary-wing aero-medical evacuation team, Kandahar; and 2ndly as the Regimental Medical Officer (RMO) to Reconstruction Task Force 3, Tarin Kowt. He has also deployed as RMO to the Force Level Logistic Asset 5, Kuwait and as MO, Resuscitation Team, Timor Leste.

David works as a Contract Health Practitioner to the ADF at Kuttabul Hlth Centre, and the ADF Ward, St Vincent’s Hospital.

He is married with a 7-yr old daughter and 3-yr old son. He has a particular interest in the complexity of ‘medico-legal-ethical’ issues encountered by health personnel on operational service.

Introduction: Was this not presented at last year’s conference? Why again? It is an issue of importance and so will be discussed once more. As the educationalist states: “Only by repeating the lesson will the idea be reinforced.”

At the 2011 AMMA Conference examples of medico/legal/ethical dilemmas during an armed conflict were discussed. The rules by which we decide how to act when confronted by such dilemmas are known as ‘International Humanitarian Law’ (IHL). IHL is as

relevant today as it has ever been. IHL is clear in relation to how health personnel should act during armed conflicts. To act counter to IHL places the health provider in an unethical and potentially unlawful position. This paper will describe attitudes to IHL by ADF personnel.

Background: IHL, or the ‘Law of Armed Conflict’, is the special branch of law governing situations of armed conflict, i.e. war. The main purpose of IHL is to mitigate the suffering caused by war. It regulates the conduct of parties to an armed conflict and provides protection to those most affected. It governs the behaviour and conduct of both combatants and non-combatants. Within IHL there are specific rules relating to health personnel.

IHL has been gradually formulated over the years and its main advocate is the International Committee of the Red Cross. IHL stems from both customary international law and international conferences and treaties, in particular the 1949 Geneva Conventions and the 1977 Additional Protocols.

The Australian parliament has incorporated IHL into domestic law. It is binding on all Australian citizens.

Comments: This presentation will draw on discussions with, and examples raised by, various ADF personnel. It will illustrate whether IHL is understood and whether it is felt to be of importance. Examples are shown where ADF personnel are confronted with real-life situations and must decide on the appropriate medico/ethical/legal route to follow. In a number of cases the answers are surprising. Opinions are expressed by a wide range of personnel: From the private medic to senior health personnel; from the infantry junior leader to the combatant commander; from logisticians to legal officers. It becomes clear that this issue raises differing opinions: Some personnel understand IHL, abide by it and fight for it. However, others demonstrate a poor knowledge of IHL and neglect it. Others argue against IHL, and some even scorn it.

The speaker will reiterate that these rules are explicit. IHL provides rules and guidance on how health personnel are to act. As a senior IHL lawyer commented: “When it is most difficult to apply/follow (IHL) ...that is when it is most needed”.

The reason why ADF personnel have differing views on IHL will be explored. Why are we not getting it right? How can we get it right? Strong leadership is required to ensure all ADF personnel act ethically. Leaders have a responsibility not just to their soldiers, sailors and airmen but to the nation to ensure that International and Australian laws are adhered to.

Conclusion: This talk will illustrate that some ADF Mbrs disregard IHL in relation to Hlth personnel. To act counter to the rules of IHL places the ADF Mbr on the 'slippery slope' of declining moral and ethical standards and into potentially dire consequences. This affects not just the individual Mbr but the Defence Force and nation as a whole.

Corresponding author: David Thompson
Email: david.thompson14@defence.gov.au

The introduction of operational automated neuro-cognitive testing in the Australian Defence Force

COL Leonard Brennan

COL Brennan joined the Australian Regular Army in 1986, he graduated from the University of Newcastle in 1989 and his posting highlights include Regimental Medical Officer 3rd Battalion Royal Australian Regiment, Officer Commanding 1st Parachute Surgical Team and Commanding Officer 1st Health Support Battalion. He also has extensive staff experience including HQ 3rd Brigade, HQ Land Command, HQ Training Command-Army, HQ Joint Operations Command and his current appointment as Director Army Health, Army HQ. COL Brennan has had operational service in Papua New Guinea, East Timor, Solomon Islands and the Middle East Area of Operations. He is a specialist general practitioner and medical administrator and an adjunct associate professor with the University of Queensland.

Research question: Can an automated neuro-cognitive test like Cogstate Sport™ (Cogstate) be a useful adjunct to the management of minor traumatic brain injury (mTBI) in the operational environment?

Background & literature review: mTBI associated with an improvised explosive device (IED) blast exposure is considered a signature injury from the recent Iraq and Afghanistan conflicts. The use of a post-blast neuro-cognitive test is not new, with the US Military Acute Concussion Assessment (MACE) being widely used. The Automated Neuropsychological Assessment Metrics (ANAMS) is routinely used pre and post deployment but is not routinely used in the acute management of mTBI on operations. In 2011 it was determined that all ADF personnel deployed or deploying to Afghanistan were to have a base-line Cogstate.

Study Design/Methods: The baseline and after-injury Cogstate Sport™ database was analysed and matched against operational blast registers and clinical records to assess compliance with ADF policy on the use of Cogstate Sport post blast.

Results: The ADF achieved base line testing in 99% personnel deployed in Afghanistan by 31 Dec 2011. In 2012, after-injury cogstate testing has been incorporated into the assessment and management of all ADF members with a significant blast exposure. Cogstate testing has demonstrated significant, reversible short term neuro-cognitive impairment and subtle post-blast symptoms that were not clinically apparent. The most current results will be presented.

Discussion: Cogstate provides facilitates a more sensitive assessment of post-blast symptoms and neuro-cognitive function compared to clinical examination or MACE alone. The key limitation, access to the internet is likely to be less of a problem on future operations.

Corresponding author: Leonard Brennan
Email: leonard.brennan@defence.gov.au

Mental Health

Suicidality in the Australian Defence Force: Results from the 2010 ADF Mental Health Prevalence and Wellbeing Dataset

Dr Miranda Van Hooff, Dr Kate Fairweather-Schmidt, Professor Alexander McFarlane, Colonel Stephanie Hodson, Mrs Helen Benassi, Dr Alan Verhagen, Mrs Nicole Steele

Dr Kate Fairweather-Schmidt completed a Science degree majoring in Psychology with honours at The Australian National University (ANU). Her doctoral research at the Centre for Mental Health Research (CMHR, ANU) addressed suicidality in a community sample at an epidemiological level, and considered the influence of variables including mental health, life

events, physical illness and employment with a special emphasis on differences across age and gender. Since then, Kate has authored and co-authored a number of papers on Suicidality. Currently a senior research fellow at the Centre for Traumatic Stress Studies (University of Adelaide), Kate is investigating predictors of suicidality in the Australian Defence Force Health and Wellbeing Survey.

Recent findings from the US have triggered concern that the rate of suicide is increasing in military populations and may surpass those in the general population.

A significant body of evidence supports a continuum of suicidal expression, where non-fatal suicidality (i.e., suicidal ideation, suicidal plans and attempts) predominantly precede future completed suicide. Thus, focusing on suicidal thoughts and behaviours provide effective opportunities to prevent suicide deaths. In Australia, the prevalence of suicidal ideation and making a suicide plan was significantly higher in the ADF compared to the Australian community, with the rate of suicidality in the ADF being more than double that in the general community. However, the ADF reported the same prevalence of suicide attempts (0.4%) in the preceding 12 months as the general community (0.3%). The psychological and physical trauma associated with combat experiences may contribute to mental disorders and place military personnel at an increased risk of suicide. This presentation will identify the risks of suicidal thoughts and behaviours (i.e., non-lethal suicidality) among ADF personnel who have been exposed to significant traumas; adverse occupational factors; who serve in different arms of the military across three main categories of rank; and finally, likelihood of suicidality for personnel who have been identified as suffering affective, anxiety and/or alcohol disorders.

Corresponding author: Miranda Van Hooff
Email: miranda.vanhooff@adelaide.edu.au

Suicide Prevention in the ADF: Up skilling the mental health provider workforce

Carole Windley

Ms Windley is the Director of Mental Health Clinical Standards and Practice in Joint Health Command. She is a psychologist with 20 years experience working in both service delivery and mental health policy. This includes a significant period in VVCS – Veterans and Veterans Counselling Service, and a secondment to DoHA working on National Mental Health Reform. Her current responsibilities in Defence involve the Early Intervention, Suicide Prevention Program; Quality Assurance and Clinical Standards in Mental Health; Alcohol, Tobacco and Other Drugs Program and the ADF Centre for Mental Health.

In 2011 Defence engaged Communicorp to undertake a training needs analysis of its mental health professional workforce and develop a Suicide Risk Assessment Training (SRAT) package. This paper will describe the development of SRAT including challenges in developing a nationally relevant training package. The training acknowledges Defence has an already skilled multi-disciplinary workforce in the assessment of suicide. However, SRAT clarifies and operationalises health policy in a competency

based format and the delivery mode ensures sustainability over the longer term.

Corresponding author: Carole Windley
Email: carole.windley@defence.gov.au

Consequences of deployment to Timor-Leste for Australian military families

Dr Annabel McGuire, Ms Catherine Runge, Dr Renee Anderson, Ms Katrina Bredhauer, Professor Annette Dobson, A/Prof Peter Nasveld

Dr Annabel McGuire joined the Centre for Military and Veterans' Health (CMVH) in January 2007 as a Research Fellow within the Deployment Health Surveillance Program. Currently, she is the Chief Investigator on the Timor-Leste Family Study. During her time with CMVH, Annabel has been co-recipient of four Defence and DVA grants, produced six peer-reviewed journal articles and more than 20 reports to Defence & DVA. Annabel has been co-author of 25 conference presentations, presenting nine of these.

The literature suggests that military families are resilient, but they are regarded as a special population because they face unique stressors. In particular, deployment separation may be associated with negative family outcomes such as poor mental health, marital dissatisfaction and child health and behavioural problems^{1, 2}. The current research explores these discrepant findings to determine the physical, mental or family health impacts Australian families experience as a result of military deployment to Timor-Leste.

The Australian Department of Veterans' Affairs funded a large cross-sectional quantitative study as part of the larger Family Study Program. Responses were collected from Australian veterans and partners (N>4,000) using validated measures of physical (SF-12), mental (K10, SF-12, PCL-C), family (FACES-IV) and child (SDQ) health outcomes associated with deployment to Timor-Leste.

Key findings on the health outcomes for partners and children in the Timor-Leste Family Study will be presented. The findings support aspects of prior research particularly from the United States and provide new insights on the health and wellbeing of Australian military families and their family dynamics. Conclusions and implications of these findings for policy, service providers and future research are addressed.

References:

De Burgh, H. T., White, C. J., Fear, N. T., & Iverson, A. C. (2011) The impact of deployment to Iraq or Afghanistan on partners and wives of military

personnel. *International Review of Psychiatry*, 23 (2): p. 192-200.

Flake, E. M., Davis, B. E., Johnson, P. L., & Middleton, L. S. (2009). The Psychosocial Effects of Deployment on Military Children. *Journal of Developmental and Behavioural Paediatrics*, 30, 271-278.

Corresponding author: Annabel McGuire
Email: a.mcguire@uq.edu.au

Telepsychiatry in the Australian Defence Force

Dr Duncan Wallace

Dr Wallace is a consultant psychiatrist at the ADF Centre for Mental Health, Mosman, NSW. He is a Commodore in the Royal Australian Naval Reserve and holds the position of Director General Health Reserves –Navy. He has undertaken several operational deployments, most recently serving in the Middle East from January to March 2012 as medical staff officer at the Australian Headquarters in the UAE.

The Australian Government's response to the Dunt Review of Mental Health services in the ADF included the establishment of a telepsychiatry hub at the ADF Centre for Mental Health. The presentation describes the development so far of a telepsychiatry network and operating policies to provide consultant psychiatrist services to garrison health and to deployed personnel on operations.

Tobacco Use and Nicotine Dependence in the Australian Defence Force: Results from the 2010 ADF Mental Health Prevalence and Wellbeing Dataset

Dr Miranda Van Hooff, Professor Alexander McFarlane, Colonel Stephanie Hodson, Mrs Helen Benassi, Dr Alan Verhagen, Mrs Nicole Steele

Dr Van Hooff is currently a research fellow at the Centre for Traumatic Stress Studies. She qualified from her honours degree in Psychology in 1998 and in 2011 was awarded the degree of Doctor of Philosophy in Medicine for her research into the longitudinal outcomes of childhood disaster exposure. Over the last 10 years she has conducted a number of large-scale longitudinal studies of traumatized populations. Miranda is an investigator on the 2010 Mental Health Prevalence and Wellbeing Study.

In military populations, cigarette smoking has been reported to increase upon enlistment, is particularly prevalent in personnel on operational deployment and has been shown to negatively affect readiness for duty, productivity and length of military service. Furthermore there is a reported association between cigarette smoking and alcohol and drug use, particularly in younger military members, which resembles the patterns in the general community. Utilising results from the 2010 ADF Mental Health Prevalence and Wellbeing Study, this presentation is the first to examine the prevalence of tobacco use and disorder in a representative sample of the ADF. Using a structured diagnostic interview (CIDI), 4.5% of the ADF (n = 2266) met criteria for ICD-10 Nicotine Dependence in the past 12 months. The prevalence of tobacco consumption across rank, service sex and deployment will be discussed.

Corresponding author: Duncan Wallace
Email: duncan.wallace2@defence.gov.au

Academic Training

Professional Development for Military Medical Officers; components towards the Diploma in Military Medicine

CDRE Jenny Firman, LEUT Brendan Peek

Jenny Firman joined the RAN in 1980 and spent 22 years in the full time RAN as a medical officer. She enjoyed a range of positions including sea postings, underwater medicine experience and Joint Health Command postings. She left full time Navy in 2002, taking up the position of Chief Medical Officer of Defence Force Recruiting and then in 2007 a position as medical adviser in the Office of Health Protection in

the Department of Health and Ageing, a role that supports national responses for health emergencies.

Throughout her career CDRE Firman has participated in general practice and is Chair of the Military Chapter of the Royal Australian College of General Practitioners.

Currently there is a limited career pathway for Defence Medical Officers wishing to remain in a generalist clinical pathway. This important group provide a significant medical cover and operational capability to the ADF. Australian Public Service (APS) and contracted medical officers in the support area also provide significant services to the ADF, often

without any specific or standardised training relevant to the military environment.

The Diploma and Certificate in Military Medicine will be a formal qualification that recognises and supports the development of advanced medical skills that are unique to the military environment, whilst acknowledging and embracing the same evidence-based, generalist medical care model that exists across Australia.

When complete, the benefits of this program will include:

- providing an “gold standard” to which military medical officers can aspire to, in terms of professional and academic excellence, whilst maintaining a generalist focus;
- providing a positive recruitment and retention tool for both full time and reserve, APS and contracted medical officers working with defence;
- providing the vehicle for greater collegiality, under the auspices of the Chapter of Military Medicine of the Royal Australian College of General Practitioners;
- enhancing the reputation of ADF medical officers to the wider professional community, the general public and to the ADF population through the recognition of the high value of military medical skills;
- providing appropriate and relevant opportunities for meaningful continuing professional development, thereby satisfying the requirements for registration by the Medical Board of Australia.

The Diploma and Certificate will have as its foundation the existing curriculum for general practice of the RACGP. It will then attempt to extend this curriculum to capture the unique context and, at times, content of military medicine. This session aims to discuss the basic content of the curriculum and review the three components of the qualification: military medical training, military medical experience and assessment.

The session will describe the development of this qualification and identify opportunities for individual involvement in the process of curriculum development to achieve this qualification and ongoing involvement and support.

Corresponding author: Jenny Firman
Email: jenny.firman@health.gov.au

Aerospace medicine as a clinical specialty – an initiative to recognize ‘specialists in aerospace medicine’

Turner JT, Smith AM, Cable GG

John Turner joined the Royal Navy as medical cadet in 1971. He graduated as a doctor from Charing Cross Hospital Medical School, University of London in 1974.

He was the Censor for Remote and Rural Trainees on the Board of Censors of the Australasian Faculty of Occupational & Environmental Medicine from 2002 and has been the Chair of the AFOEM Teaching and Learning Committee since 2009.

After an enjoyable year (2005-6) as the Chief of Operations at the Centre for Military and Veterans' Health (CMVH), University of Queensland he had the good fortune to be able to return to pure Aviation Medicine as the Commanding Officer RAAF IAM (2008- 2010). He is currently enjoying being back at work as a reservist with the Army in Townsville.

He feels particularly honoured to have been asked earlier this year to take on the role of Inaugural President of the Australasian College of Aerospace Medicine (ACAsM).

Aerospace medicine is seen as a clinical specialty in many parts of the world; however, in Australia and New Zealand it has been viewed for many years as a ‘special interest group’ of general practitioners, occupational physicians, and military doctors. Within this paradigm, interest to pursue a career in aerospace medicine has diminished - to the point it has become difficult to identify people with the knowledge and experience to fill key functions in aerospace medicine. The Australasian College of Aerospace Medicine was established to provide an academic structure to support and develop those doctors in Australia and New Zealand who practice aerospace medicine at a high level of expertise, and to support and train junior doctors who aspire to a career in aerospace medicine - with an aspirational goal of aerospace recognized as a clinical specialty by the wider medical community. This presentation will outline the College's model for seeking specialist recognition, and describe the way it is achieving these goals. Not only will this presentation interest prospective specialists in aerospace medicine, but it will also contribute to a discussion of how to recognise as 'specialists' those people who practice niche areas of military medicine to a high level of expertise - especially in disciplines where there is no clear civilian counterpart.

Corresponding author: Adrian Smith
Email: adrian@pegasusaeromed.com

The ADF Chair of Military Medicine and Surgery

LTCOL Michael C. Reade

After clinical training in Sydney and Melbourne, a doctorate from Oxford and an MPH from the University of Pittsburgh, Michael returned to Melbourne as an Associate Professor of Intensive Care Medicine in 2007. In 2011, after 22 years in the Army Reserve including deployments to the Balkans, Timor, the Solomon Islands and Afghanistan, he was appointed to a full-time Defence position as the Professor in Military Medicine and Surgery at the University of Queensland. He is developing a research programme at the interface of civilian and military trauma, including management of acute cognitive impairment, traumatic coagulopathy, and trauma systems design.

In 2011, the Australian Defence Force built on existing health research strengths at the Australian Malaria Research Institute and the Centre for Military and Veterans' Health by establishing the ADF Chair of Military Medicine and Surgery within the Burns, Trauma and Critical Care Research Centre at the University of Queensland. By adding trauma medicine and surgery to research in tropical medicine and occupational and public health, the ADF now supports a research hub in southern Queensland relevant to all aspects of military medicine.

The tasks of the ADF Chair are to lead a programme of original research relevant to both military and civilian trauma, and to guide the implementation of modern trauma practice in the ADF. Research projects currently underway include:

- evaluation of cryopreserved blood products and clotting factor concentrates, in collaboration with the Australian Red Cross Blood Service, the Australian and New Zealand Intensive Care Research Centre, and the Queensland University of Technology Medical Engineering Research Facility;
- clinical trials of novel approaches to the management of traumatic brain injury and acute cognitive impairment, in collaboration with the Australian and New Zealand Intensive Care Society Clinical Trials Group;
- clinical evaluation of rapid molecular diagnostic testing for early evidence of infection, in collaboration with the University of Western Australia; and
- evaluation of various trauma systems designs, in collaboration with the National Trauma Research Institute and the Queensland Trauma Registry.

Further studies in the fields of inflammatory-modulating trauma fluid resuscitation and blast and blunt lung injury are planned, in collaboration with

the United States Institute of Surgical Research and the UK Defence Science and Technology Laboratory Porton Down and the Royal Centre for Defence Medicine. ADF officers engaged in civilian trauma research around Australia are invited to consider the potential funding and impact benefits of linking their work to the ADF programme.

The ADF Chair also has input into the development of the deployed ADF trauma system. This includes:

- oversight of the academic aspects of the fulltime ADF procedural specialty registrar training programme, including liaison with the specialty colleges;
- better integration of the training provided in clinical short courses, such as the EMST;
- development of a process of training and credentialing military surgical teams prior to deployment;
- development of ADF clinical practice guidelines relevant to trauma care, including those of the Australian Resuscitation Council;
- implementation of a process of clinical audit and review of the work of deployed medical teams; and
- development of an ADF trauma registry.

In time, it is hoped that the concentration of ADF clinical trauma practice, training and research within the one institution – perhaps termed an 'Australian Centre for Defence Medicine' – will facilitate the efficiency, corporate identity and effectiveness that typifies similar leading clinical institutions worldwide.

Corresponding author: Michael Reade

Email: m.reade@uq.edu.au

Masters of Military Medicine – Enhancing Medical Officer Education & Training

COL Richard Mallet, MAJ Carol Waldeck

COL Richard Mallet is the Command Health Officer for Headquarters Force Command. COL Mallet and his team are responsible to the Commander Forces Command for ensuring that FORCOMD has a robust and relevant deployable Combat Health capability to support both deployed and non-deployed forces.

A measured and more defined response to education and training for Army Medical Officers is required following the endorsement of the Combat Health Operating System as a key component of the Army Health Force Modernisation. This includes the ongoing development of supporting sub-systems and associated capabilities. The effective generation of the Combat Health Operating System requires

adherence to detailed and iterative clinical education and training regimes within foundation war-fighting and mission specific programs. Future education and training pathways for Army Medical Officers must be specifically targeted towards combat health capability needs, whilst retaining the flexibility to be relevant across the spectrum of conflict. Effective health training and education underpins the operation of Army combat health capability in complex and challenging environments. The readiness of combat health personnel is dependent on the successful synthesis of individual and collective education and training. This must be reinforced by current and relevant clinical practice.

Education and training for Army Medical Officers is currently not optimised to deliver the specific needs of Army's operational health requirements. Current education and training pathways have not been developed specifically to meet the needs of Army Medical Officers in complex and austere environments. It is also recognised that relevant

clinical skills for Army Medical Officers are highly erodible and historically difficult to maintain. This vulnerability applies to pre-hospital emergency care, primary health care or high-end deployed hospital care. This is due in part to the limited ability of Army Medical Officers to fully utilise their skills in non-deployed military environments. For Army Medical Officers, education and training consists of a mix of professional clinical training, in-house military medical training and medical education & training external to Army. The drivers for the Masters of Military Medicine have been made in recognition of the current context and lessons-learned on operations over the past decade. In addition, the Masters of Military Medicine serves to mitigate the vulnerability in the current education and training provided to Army Medical Officers and aims to set the conditions for Army's future combat health capability requirements.

Corresponding author: Richard Mallet
Email: richard.mallet@defence.gov.au

Miscellaneous

The Effect of Centre-Based Counselling for Veterans and Veterans' Families on Long Term Mental Health Outcomes

David Forbes, O'Donnell, ML., Varker, T., Perry, D.

David is the Director of the Australian Centre for Posttraumatic Mental Health (ACPMH) and Professor, Department of Psychiatry, University of Melbourne. He is a clinical psychologist with many years' experience in the assessment and treatment of mental health problems following trauma.. He was the Chair of the working party for the Australian Guidelines for the Treatment of Adults with Acute Stress Disorder and Posttraumatic Stress Disorder. He is on the Editorial Board of key international journals and has published widely in the area of posttraumatic stress in the international literature

Objective: The Veterans and Veterans Families Counselling Services (VVCS) was established by the Australian government and plays a pivotal role in providing mental health services to veterans and their families. This research explores the impact of centre-based psychological counselling upon depression, anxiety, stress and alcohol use severity.

Methods: We invited a stratified sample of VVCS clients to participate in this study. Data was collected on intake to the program, at the fifth counselling session, and 12 months after the commencement of counselling. Repeated measures General Linear

Model (GLM) analyses were conducted to examine the impact of centre-based counselling upon depression, anxiety, stress and alcohol severity over time. Secondary analyses were conducted to investigate the role of gender, client status and era of service on changes in symptoms across time. These analyses allowed us to compare pre- and post-counselling client outcome data.

Results: VVCS centre-based counselling resulted in a significant reduction in depression, anxiety, stress and alcohol use severity by the end of counselling, and these improvements were maintained over the next 12 months. Females responded particularly well to centre-based counselling, and contemporary veterans overall had lower levels of symptom severity than older veterans over time.

Conclusions: VVCS centre based counselling successfully reduced depression, anxiety, stress and alcohol use symptom severity of veterans and their families. Contemporary veteran and females responded particularly well to VVCS centre based counselling.

Corresponding author: David Forbes
Email: dforbes@unimelb.edu.au

The Great War and the Brisbane Hospital

Assoc. Prof. Cliff Pollard

General surgeon; recent retired as Director of Trauma Services RBWH; currently has a clinical governance/audit post in Retrieval Services Q'ld; Chair of the Statewide Trauma Clinical Network; deployed to Bougainville and East Timor with the RAAMC.

When the Great War broke out in 1914, the implications for the Brisbane Hospital were immediate and would continue for the duration and beyond of the conflict. Staff resigned from the Hospital at the beginning of the War and throughout to enlist. At one stage, the Hospital had no resident medical officers. Grace Wilson, the matron in August, 1914, would enlist immediately and would go on to achieve the highest nursing post in the Australian Army. Others would provide outstanding service with due recognition and subsequently have impressive post-war nursing careers. One medical officer would take his own life after the Dardanelles campaign. John Barr McLean would return to his post as superintendent until retirement in 1933. This paper will examine in part the impact on the Brisbane Hospital, and also record some of the events in the lives of this extraordinary group of people who enlisted from it.

Corresponding author: Cliff Pollard

Email: Cliff_pollard@health.qld.gov.au

More than Malaria. The contribution of Australian Army Doctors to the science of Tropical Medicine in the two World Wars.

Professor Geoffrey Quail

Geoff Quail has a long association with the RAAF since serving as a national serviceman till retirement as consultant in medical education and tropical medicine in the Specialist Reserve. He was appointed lecturer in the Department of Oral Medicine in 1961 and has held continuous appointments at Melbourne and Monash Universities since that time. He is a councillor of the Australasian College of Tropical Medicine and was a Centenary Lecturer for the Royal Society of Tropical Medicine and has received the AMMA Research Award on two occasions.

Geoff read modern history at Oxford and is currently writing a book on the Contribution of the Australian Military to the Science of Tropical Medicine.

Much has been written about the impact of malaria

on the allied forces in the Pacific and Burma campaigns of the Second World War but there is little documentation of the other diseases which caused serious morbidity and greatly reduced fitness in troops serving in the tropical and sub-tropical theatres during both world wars. During these times of conflict Australian military doctors and scientists greatly advanced knowledge of these diseases which included scrub typhus, bacillary and amoebic dysentery, schistosomiasis and dengue fever. Their skill and dedication resulted in a significant reduction in morbidity of these conditions and so greatly improved the health of serving members. This translated to improved health of people living in the tropics after hostilities ceased. This paper will describe the impact of these diseases at Gallipoli, the Sinai, Burma and New Guinea and the significant advances which resulted from Australian military doctors and scientists' scientific work.

Corresponding author: Geoffrey Quail

Email: geoffrey.quail@med.monash.edu.au

The Case of the Pelvic Digit

Peter Hurly

WGDCR Peter Hurly is currently the Director of Air Force Medicine for the RNZAF. He has been a member of St John Ambulance since 1964 and was involved in ambulance work and training. He trained as a pharmacist in South Africa and studied medicine obtaining his MBChB in 1983. He worked in hospital medicine and Accident and Emergency. He was a member of the South African Military Medical Service and saw active service in South Africa. On moving to New Zealand, he joined the Royal New Zealand Army Medical Corps. He then moved to general practice in Palmerston North and took up a Reservist position with the RNZAF. He obtained a Diploma in Aviation Medicine and a certificate in Air Retrieval medicine, subsequently progressing to a Masters in Aviation medicine. He moved back into full time military medicine in 2002 and became the Director of Air Force Medicine in 2004. He is due to retire from the Regular Force at the end of this year.

A case presented of an unusual congenital condition in a service person, a discussion of the presentation and management and a literature search of the condition followed by an indication of treatment options and prognosis.

Corresponding author: Peter Hurly

Email: peter.hurly@nzdf.mil.nz

Operational Health

Vom militarisches sanitats leistungen: On military health services

AIRCDRE Warren Harrex, COL John Turner, CMDR Neil Westphalen

John Turner joined the Royal Navy as medical cadet in 1971. He graduated as a doctor from Charing Cross Hospital Medical School, University of London in 1974.

He was the Censor for Remote and Rural Trainees on the Board of Censors of the Australasian Faculty of Occupational & Environmental Medicine from 2002 and has been the Chair of the AFOEM Teaching and Learning Committee since 2009.

After an enjoyable year (2005-6) as the Chief of Operations at the Centre for Military and Veterans' Health (CMVH), University of Queensland he had the good fortune to be able to return to pure Aviation Medicine as the Commanding Officer RAAF IAM (2008-2010). He is currently enjoying being back at work as a reservist with the Army in Townsville.

He feels particularly honoured to have been asked earlier this year to take on the role of Inaugural President of the Australasian College of Aerospace Medicine (ACAsM).

Traumatic brain injury, post-traumatic stress disorder and traumatic amputations have drawn considerable attention from political and military leaders, veteran's groups and the media. Such injuries have become the 'signature wounds' of the Iraq and Afghanistan wars. Furthermore, comparable attention has often been expended on medical conditions that can be characterised by their novelty, severity and/or lack of ascribable causation.

However, the focus on these short-term military health imperatives has diverted attention from the long-term strategic principles of military health care, in the context where information on the functions and roles of military health services is at best limited, or at worst non-existent.

This in turn supports the premise that military health services are perceived and externally, simply as uniformed treatment services for use primarily in operational circumstances. It is this perception, combined with the focus on short-term military health imperatives, that currently drive how many Western military health services are organised and resourced.

This presentation will describe the functions and roles of national military health services at the strategic level, where they are applicable irrespective

of the size of the military services that they support.

*Corresponding author: Neil Westphalen
Email: neil.westphalen1@defence.gov.au*

Pilot Australian Defence Force Military Surgical Team at Royal Brisbane and Women's Hospital 2012

Dr Amanda Dines, CMDR Alison Thomas

The ADF Military Surgical Team will comprise five specialist positions in: emergency medicine, general surgery, orthopaedics, anaesthesia; and intensive care. Each Member of the ADF Military Surgical Team will be a member of the Reserve Force and will be released from their duties at the RBWH to undertake Defence Service as required by the ADF for up to 16 weeks per year.

At the RBWH, members of the ADF Military Surgical Team will develop their clinical skills as individuals and as a team, with a clear focus on a trauma based approach to clinical practice. The ADF Military Surgical Team will be integrated within the RBWH clinical team; be involved in the delivery of training to ADF personnel on secondment to the Hospital; and be involved in research and teaching.

During military service, the ADF MST will contribute to and enhance the ADF's health support capability on military and humanitarian operations and exercises.

The presentation will discuss the implementation and key outcomes of the pilot project in 2012.

*Corresponding author: Amanda Dines
Email: adines2@me.com*

Providing Effective Forward Surgery in Manoeuvre Warfighting – A Command Perspective

Mark Elliott

Major Mark Elliott commanded 23 Air Assault Medical Squadron during the 2003 invasion of Iraq whilst in the British Army. The Squadron provided Role 1 and 2 (enhanced) forward surgery to manoeuvre Battlegroups during all phases of the operation. He has a wide range of operational experience covering peace support, stability and counter insurgency operations and has deployed twice on manoeuvre warfighting operations.

He is a lateral recruit to the ADF and has worked in the Land Warfare Development Centre. His current role is in Health Plans in HQ Northern Command.

The Invasion of Iraq in 2003 was the last time that formation manoeuvre warfighting occurred in the British Army. Current operations see health support occurring within a framework of the Joint Trauma system and that system currently supporting a mature operational environment. The mature operational environment is generally based on a framework of forward operating bases and support bases with dedicated Aeromedical evacuation support.

In formation manoeuvre warfighting you are faced with a completely different operational environment. The ability to be able to provide effective support at the Battlegroup (BG) and Brigade level relies on being embedded in the planning phase with the BG HQ, a comprehensive command and communication suite to be able to talk to both ground and air assets, mutual support from other health units and a medical facility that is adaptive, agile and robust enough to be able to deploy and re-deploy for a range of mission types. Personnel training and skillsets have to be matched to the mission and equipment available.

Further there are a range of constraints and restrictions that have to be planned and factored for including an uncertain casualty estimate and poorly defined "medical rules of engagement", "no fly" periods for Aeromedical evacuation due to availability, threat level, weather or offensive support, health logistics issues, protection, mobility and environmental factors both internal and external.

23 Air Assault Medical Squadron centre of gravity was the Air Assault surgical group (AASG) providing Role 2 (Enhanced) Forward Surgery (including Damage control Surgery) with Medical Sections deployed out providing Role 1 support. The AASG shape and form was the result of a distillation of over 20 years of operational experience which resulted in a facility that was adaptive and robust enough to deploy on a range of air platforms (including parachute) and by a wide variety of land assets.

This paper will outline the issues faced in providing forward surgical support in this environment and discuss in detail what worked and lessons learnt.

Corresponding author: Mark Elliott

Email: mark.elliott6@defence.gov.au

Controversies in trauma resuscitation: plasma-to-red cell ratios, platelets, tranexamic acid, and hypotensive resuscitation

LTCOL Michael C. Reade

After clinical training in Sydney and Melbourne, a doctorate from Oxford and an MPH from the University

of Pittsburgh, Michael returned to Melbourne as an Associate Professor of Intensive Care Medicine in 2007. In 2011, after 22 years in the Army Reserve including deployments to the Balkans, Timor, the Solomon Islands and Afghanistan, he was appointed to a full-time Defence position as the Professor in Military Medicine and Surgery at the University of Queensland. He is developing a research programme at the interface of civilian and military trauma, including management of acute cognitive impairment, traumatic coagulopathy, and trauma systems design.

Modern research has questioned many time-honoured practices in trauma resuscitation, and introduced promising new technologies. However, evidence supporting these developments seems prone to unwarranted extrapolation by enthusiastic clinicians.

The British and US militaries implement a policy of 1:1 fresh frozen plasma: packed red blood cell ratio for transfusion in patients anticipated to require massive transfusion (>10 units in 24 hours) due to trauma. However, the Australian National Blood Authority in its 2010 guideline recommends only that a ratio of 1:2 be considered, and the American Association of Blood Banks 2010 guideline explicitly makes no recommendation for or against a ratio of greater than 1:3. By March 2012, 26 observational studies enrolling 6655 patients (Ho AM et al., *Anesthesiology*, 2010) had addressed this question. Uncertainty and lack of agreement in guidelines relates to possible survivor bias in many of these studies. There is undoubtedly also a conflict between clinicians tempted by an intuitively appealing concept, and custodians of the blood supply who see potential broader implications. The Australian Defence Force currently makes no recommendation, though anecdotally a policy of 1:1 transfusion has been practised by the Australian Defence Force Medical Treatment Facilities in the Middle East.

A high ratio of platelets to other blood components transfused has also been found in retrospective data to be associated with improved outcome. However, the evidence is even less sound than for plasma, and the availability of liquid-stored platelets (with a shelf-life of only five days) limits the applicability of this therapy. Cryopreserved platelets overcome problems of supply, and are possibly more clinically effective. ADF personnel have received cryopreserved platelets during recent deployments, and the Dutch armed services have more than six years' experience with this technology. However, human clinical trial evidence is limited to a single study of 73 patients.

The recent CRASH-2 trial (CRASH-2 collaborators, *Lancet*, 2010) found mortality reduced by tranexamic acid if given within three hours of injury. However, a

subsequent analysis found death due to bleeding significantly increased (from 3.1% to 4.4%) if tranexamic acid was given later than 3 hours. Furthermore, the trial was conducted principally in developing countries, with less access to plasma transfusion and with less availability of modern trauma surgery and intensive care. The Australian Defence Force currently makes no recommendation regarding tranexamic acid, and Australian civilian trauma services have different approaches.

Hypotensive resuscitation has been central to trauma management since the landmark study of Bickell (NEJM, 2004). However, the circumstances of this study are seldom relevant to military trauma, and there is increasing concern that Acute Traumatic Coagulopathy, and indeed mortality, is worse with the prolonged application of this approach.

This review summarises the available evidence for and against the interventions listed, and makes interim recommendations for practice in anticipation of the results of the planned upcoming (hopefully definitive) trials. The ADF / Australian Red Cross Blood Service programme of research into cryopreserved platelets and factor concentrates will be a major influence on future trauma practice, both military and civilian.

Corresponding author: Michael Reade

Email: m.reade@uq.edu.au

Frozen blood products: a solution for deployed health facilities?

CMDR Anthony Holley

Anthony is a dual qualified intensivist and emergency physician working at Royal Brisbane and Women's Hospital as a senior staff specialist. He is a senior lecturer with the University of Queensland Medical School and Chairman of Australia New Zealand Intensive Care Society, Queensland. Anthony was the recipient of the Matthew Davey Award that facilitated study at the Netherlands Military Blood Bank. He serves as a representative for the National Blood Authority Critical Care Expert Group. Anthony has deployed on a number of occasions including to

Angola, Bougainville, East Timor, The South Pacific, The Persian Gulf and most recently to Afghanistan.

Haemorrhage is a potentially reversible cause of trauma deaths. The concept of haemostatic resuscitation, characterised by transfusion of blood products in an immediate and sustained fashion is well established. Early transfusion with red blood cells, platelets and plasma in a 1:1:1 ratio appears beneficial. This paradigm demands products be readily available in austere environments. Investigation into synthetic products has failed to provide a viable alternative. Refrigerated liquid products are limited by short shelf lives, while fresh, warm whole blood has significant draw backs. Deep freezing is able to substantially extend the shelf life of blood products required for resuscitation, facilitating practical resupply.

Product	St	Storage shelf Life of product
	Standard Liquid Blood	Dep Frozen Blood
Red Blood Cells	42 days at 4oC	10 years at -80oC
Platelets	5 days at 22oC	2 years at -80oC
Fresh Frozen plasma	1 years at -30oC	7+ years at -80oC

Deep frozen blood product production requires cryopreservation with glycerol in the case of erythrocytes and dimethyl sulphoxide for the preservation of platelets. These agents protect the biological elements from the destructive potential of deep freezing. Prior to transfusion the products must be washed/prepared including removal of cryopreservatives. The development of a deep frozen blood product supply could substantially enhance the ability to provide high quality critical care for injured service personnel on deployment or civilians in remote centres. This presentation will describe the processes, training, Dutch experience and potential pitfalls of adopting "deep freezing" technology.

Corresponding author: Anthony Holley

Email: anthony_holley@health.qld.gov.au

Rehabilitation

Rehabilitation and Recovery of Military Personnel with Serious Wounds, Injuries and Illnesses

Jim Porteous

The Simpson Assistance Program (SAP) aims to support and enhance the efforts of Defence to reduce the impact of serious injury or illness on Australian

Defence Force (ADF) personnel. Defence currently has a very successful occupational rehabilitation program but acknowledges that improvements can still be made. SAP will assist and encourage ADF personnel (with the support of their families) to return to the ADF workplace in either similar or new capacities through the provision of a comprehensive

approach to tailored recovery programs which support the individual needs of wounded ADF personnel and their families.

SAP is consistent with current efforts to improve family engagement, service integration and seamless transition.

There are currently 11 work streams within the SAP which contribute to rehabilitation excellence. The focus of this presentation will be on the Intensive Recovery Program, meaningful engagement and rehabilitation research projects.

Corresponding author: Jim Porteous
Email: jim.porteous@defence.gov.au

Musculoskeletal conditions and psychological comorbidities in Australian Gulf War veterans

Dr Helen Kelsall, Dr Dean McKenzie, Prof Andrew Forbes, Ms Mina Roberts, Prof Malcolm Sim

Dr Helen Kelsall is a Senior Research Fellow and public health physician at the Department of Epidemiology and Preventive Medicine, Monash University. She is an Investigator on the current study of the health of Australian Gulf War veterans and military comparison group being undertaken by Monash University and was an Investigator and undertook her PhD on the 2000-02 baseline study. Her particular interests include physical health assessment in veterans and the relationship between physical health, psychological health and exposures in veteran and military populations. Research interests include public and occupational health and continuing professional development.

Introduction: Musculoskeletal conditions are common causes of pain and disability and have been found by community studies to co-occur (be comorbid with) psychological conditions such as depression and anxiety.* Little is known about such musculoskeletal comorbidity in defence and veteran populations, however.

Aim: The aim is to investigate the comorbidity of musculoskeletal disorders and psychological disorders in Australian Gulf War veterans compared with a military comparison group.

Methods: In 2000-2002, Australian Gulf War veterans and military comparison group members participated in a study that included a comprehensive postal questionnaire, which asked about demographic and military service details, reported doctor diagnosed medical conditions including arthritis or rheumatism, joint problems, back or neck problems, fibrositis or fibromyalgia or soft tissue injuries; and

whether they had been treated by a doctor in the past year. Medical practitioners rated self-reported medical conditions as probable diagnoses, possible, unlikely or non-medical as part of a face-to-face interview. We studied musculoskeletal conditions that were diagnosed as probable and had been treated in the past 12 months. The Composite International Diagnostic Interview (CIDI) assessed current depression, posttraumatic stress disorder (PTSD) and alcohol abuse or dependence based on DSM IV criteria based on symptomatology occurring within the previous 12 months. Due to small numbers of female Gulf War veterans, these analyses were limited to 1424 male veterans and 1548 male comparison group subjects. Logistic regression was used to investigate the relationship between musculoskeletal conditions and Gulf War service or psychological comorbidity, adjusting for potential confounding variables such as age and rank.

Results: The proportion of Gulf War veterans and comparison group members who reported any doctor diagnosed musculoskeletal condition that had been treated in the past year was similar in both groups (12.6 vs 11.4%, OR 1.11; 95% CI 0.88-1.41). For arthritis or rheumatism the findings were 3.6 vs 3.5% (OR 1.09; 0.73-1.63); for joint problems were 11.9 vs 10.8% (OR 1.12; 0.88-1.41); and soft tissue disorders, both 0.4% (OR 1.10; 0.35-3.40). A slightly higher proportion of Gulf War veterans reported having treated back or neck problems (16.2 vs 13.9%, OR 1.22; 0.99-1.51) but the difference was not statistically significant. Arthritis or rheumatism was associated with comorbidity of depression in Gulf War veterans (OR 3.42; 1.64-7.14) and comparison group (OR 4.05; 1.84-8.95); with PTSD in Gulf War veterans (OR 2.89; 1.22-6.86) and comparison group (OR 3.71; 1.03-13.36) and with alcohol abuse or dependence in comparison group subjects only (OR 2.77; 1.31-5.84).

Conclusions: The occurrence of musculoskeletal disorders is similar in Gulf War veterans and comparison group. Arthritis or rheumatism is associated with increased psychological comorbidity in both study groups, consistent with community findings. Comorbidity of these conditions has implications for treatment and management and so should be considered during assessment.

*AIHW 2010. When musculoskeletal conditions and mental disorders occur together. Cat. No. AUS 129. Canberra: AIHW

Corresponding author: Dr Helen Kelsall
Email: helen.kelsall@monash.edu

Mental Health

The health and wellbeing of ADF reservists: A review of the literature

Samantha Cromptvoets, PhD

Dr Samantha Cromptvoets, BSC (hons) Melb PhD ANU, is a Sociologist and Research Fellow in the ANU Medical school. Dr Cromptvoets is Chief Investigator on a number of DVA and Defence funded projects examining issues of gender, workforce, identity, culture, health and wellbeing. Her current projects include examining the health and wellbeing of female veterans, and the history, identity and wellbeing of ADF reservists. Other research in particular includes an analysis of the free text responses in the recent ADF census and reserve attitude survey, compiling a history of the Defence Reserves Association and analysing the marketing and recruitment campaigns aimed at Reserves since Federation.

This paper presents findings from a systematic review of the academic and government literature on Reserves. The findings highlight a set of issues for Reservists that at one level exist as structural ADF personnel matters, for example conditions of service, retention, incentives, and employer and family support. However all of these have a direct impact on reservist health and wellbeing, in particular their mental health.

The main themes that emerged from the review included the history and role of Reserves internationally and in Australia, descriptions of and challenges to Reservist identity and the context of military culture, commitment and engagement, deployment and its effects on mental health, civilian employers and civilian employment, job satisfaction and Reservist spouses and families.

In conclusion, considerable gaps exist in our knowledge of the health and wellbeing of ADF Reservists. To date, most research on Australian reservists focuses on capability. Rather than dismissing this literature as irrelevant to health this review has examined the findings and found that significant factors contributing to reservist health and wellbeing are visible as issues of 'commitment' and 'retention'.

*Corresponding author: Samantha Cromptvoets
Email: samantha.cromptvoets@anu.edu.au*

The Wellbeing Toolbox: Findings of the evaluation of an online mental health and wellbeing resource

Delyth Lloyd, Chris Clarke, John O'Connor, Kym Connolly, Jane Nursey, Andrea Phelps

Kym Connolly is the Director of Mental Health Programs at the Department of Veterans' Affairs. The Department supports a wide range of mental health services, including counselling and referral services for veterans, war widow/ers, serving members, former defence force members and their families. Ms Connolly leads the delivery of client and staff mental health literacy, allied mental health provider engagement and training, and suicide prevention programs.

Jane joined ACPMH in 2010 as a Senior Clinical Specialist. She is a Clinical Neuropsychologist and has worked in public acute and mental health services across Melbourne for 18 years, including in Child and Adolescent, Adult and Aged Care services. She has developed psycho-education programs on the neuropsychology of PTSD, delivered group and individual based treatments to traumatised clients and coordinated a treatment service for bushfire survivors following the Victorian bushfires in February 2009. At ACPMH Jane is involved in policy and service development projects as well as education and training programs.

The Wellbeing Toolbox is a self-help online resource for veterans and other former serving members and their families. The primary goal of the Wellbeing Toolbox is to provide an accessible resource for those with wellbeing concerns, who are not currently engaged in other mental health services. The Wellbeing Toolbox comprises six modules, based on Skills for Psychological Recovery (SPR; National Center for PTSD and National Child Traumatic Stress Network, 2007). The modules are: solving problems; building support; helpful thinking; getting active; keeping calm; and sleeping better. A 12 month trial of the Wellbeing Toolbox commenced in February 2011. The evaluation was designed to answer the following key questions: Who is using the Wellbeing Toolbox?; How much are the different parts of the site being used?; Is it helping users to address their wellbeing problems?; Is it meeting the needs of contemporary veterans?; How acceptable and accessible is the Wellbeing toolbox?; How could the user experience be improved? Data for the evaluation was drawn from Google Analytics website tracking data, website feedback, a brief on-site survey and

interviews with users. This presentation will present the key findings and lessons learned from the trial.

Corresponding author: Kim Connolly

Email: kym.connolly@dva.gov.au

Help Seeking, Stigma and Barriers to Receiving Care in the ADF: The mediating role of mental health disorder

Hodson S, Van Hooff M, McFarlane A, Benassi H, Verhagen A, Steele, N

Colonel Stephanie. Hodson graduated from James Cook University in Townsville with a BPsych (Hons) in 1990 and joined the Army in August 1991. She had had a range of posting across Australia including recruiting, research and counselling duties. LTCOL Hodson completed her doctoral studies investigating the longitudinal psychological effects of operational deployment to Rwanda in 2002 and in 2003 completed Command and Staff College. In 2006 she assumed command of the 1st Psychology Unit and was responsible for all land base psychology support to ADF operations. While CO 1 Psych she had the opportunity to deploy to both the Middle East Area of Operations and Timor L'Este. For her work during this posting she was awarded the Conspicuous Service Cross in the 2009 Australia Day Honours List, "For outstanding achievement as the Commanding Officer, 1st Psychology Unit". She is currently psychologist adviser at Department of Veterans' Affairs

Given the prevalence of mental disorders in the ADF, we must ask why so few military personnel receive care. Research indicates that two main factors contribute to the low uptake of mental health care: the fear of stigma and perceived barriers to care. The degree to which public stigma is internalised will influence the extent that personnel feel able to access support for mental health issues. This is particularly pertinent to those with a mental health problem as it is often only when a person attempts to seek help, do they feel the stigmatization and judgment for doing so. Utilising results from the 2010 ADF Mental Health Prevalence and Wellbeing Study, this presentation will examine the relationship between current psychiatric disorder, help seeking, stigma and barriers to care. The implications of these findings for training and service provision within the ADF will be discussed.

Corresponding author: Miranda Van Hooff

Email: miranda.vanhooff@adelaide.edu.au

Group Cognitive Behavioural Therapy Treatment for Service Related Posttraumatic Stress Disorder: Effectiveness, Sustainability and Repeatability.

Andrew Khoo, Michael T. Dent, Tian P.S. Oei

Dr Andrew Khoo completed his undergraduate medical degree from the University of Queensland in 1994 and received his fellowship from the Royal Australian and New Zealand College of Psychiatrists in 2002, receiving a College medal for his final year dissertation on Post Traumatic Stress Disorder. He presently divides his time between private practice and being the Clinical Director of the Toowong Private Hospital Group Therapy Day Programs. Given his clinical practice in the private sector, his specific interests include the treatment of mood and anxiety disorders, particularly psychopharmacology and group therapy. He is a Senior Lecturer for the University of Queensland Medical School and what time he finds for academic work is currently spent compiling and publishing papers on PTSD, CBT and group therapy.

Objective: To assess 12 month outcomes of Australian combat veterans with posttraumatic stress disorder (PTSD) who participated in a six week group based CBT program at the Toowong Private Hospital. The study population included 496 consecutive admissions to the program between 1999 and 2008.

Method: Self report measures of PTSD, depression, anxiety, anger, alcohol use, relationship satisfaction and quality of life parameters were collected at intake and 3, 6 and 12 months post intake.

Results: Statistically significant and sustained improvements were noted in 12 month outcome measures for PTSD, depression, anxiety, alcohol use, anger, and quality of life. PTSD symptom reduction occurred consistently each year for nine years and exhibited an aggregated effect size of 0.68. Maximal symptom change occurred during the intensive phase of treatment (ie over the first 12 weeks) on all symptom scales and for quality of life scores. Using calculated reliable change data for the PCL, 6 out of 10 participants were found to have had a clinically reliable improvement in their PTSD symptoms sustained at 12 months with 2 out of ten no longer meeting criteria for PTSD.

Conclusions: This naturalistic research demonstrates that treatment administered under clinical conditions produces equivalent magnitudes of positive change in terms of PTSD symptoms when compared with existing efficacy data in individual and group treatments. Further these symptomatic gains are sustainable and consistently reproducible. The benefits noted from group therapy were seen as

independent of whether or not individual treatment was in place.

Corresponding author: Andrew Khoo
Email: akhoo@toowongprivatehospital.com.au

Operational Health

Environmental Health Provision in Humanitarian Crisis

Glenn Keys, James Williams

James Williams is the Project Manager - Environmental Health for Aspen Medical, a private medical solutions company based in Canberra. James is a degree-qualified environmental health consultant and has experience managing health service contracts for various Australian Government assistance missions including the Regional Assistance Mission Solomon Islands (RAMSI), International Stabilisation Force (ISF) East Timor and recently consulted for AusAID in Papua New Guinea. James has a demonstrated operational knowledge of public and environmental health challenges facing the Pacific region.

When disaster strikes, whether through natural or anthropogenic causes, the first response that often comes to mind is a classical medical approach to treat the sick or injured. However, environmental health officers can play a very large part in this first response by utilising their knowledge in potable water, food and sanitation, playing an integral role in preventing the spread of disease and limiting the duration of the emergency. This session describes James' recent experience as the environmental health consultant for AusAID during a cholera outbreak in South Fly and Middle Fly districts of Western Province, Papua New Guinea. It aims to provide an insight into Australia's northern border region, explores some of the challenges in providing technical assistance to an environmental and public health emergency in a developing country context. Although this emergency was not caused by a natural disaster, it is an example of what occurs when public health infrastructure breaks down and/or is inadequate.

Corresponding author: Glenn Keys
Email: gkeys@aspenmedical.com.au

Working Overseas - Medical Assistance and Evacuation

Dr Mike Broady, International SOS - Regional Medical Director - Assistance

Mike Broady directs the International SOS Australasian alarm centre and assistance services.

The Australasian region is known for its remote and challenging locations coupled with highly variable

local medical services. Dr Broady directs the team of doctors, nurses, security and logistics team to ensure the safety and security of our clients in these environments.

In addition to managing the Australasian region, Dr Broady's team support our 26 other alarm centres to manage health and security risk across the world.

Mike has more than 16 years of broad medical experience and has served as a Community Trust medical director in the UK, and has advised local government and the Strategic Health Authority during the Swine Flu pandemic.

Working Overseas - Medical Assistance and Evacuation

Description of Material: Over the past decade, governmental and non-governmental organisations have more employees working, living and traveling in previously remote destinations and austere environments. Operating in these regions can pose significant health challenges specific to that part of the world. Unfortunately, the medical infrastructure may be significantly limited across many regions globally and individuals needing healthcare abroad in these environments may require medical assistance or evacuation.

Application: Medical assistance ranges from medical advice to emergency medical transportation to save life or limb. Aeromedical movement can take a variety of forms ranging from air ambulance evacuations to medically escorted transportation on a commercial airliner. Each patient needs to be assessed on a case-by-case basis taking into account a variety of factors to include barometric effects, oxygen limitations, and mobility. An understanding of the complexities and limitations of international aeromedical evacuations allows governmental and non-governmental organisations to be better prepared in the event of an emergency.

Results, Observations, and Conclusions: Original analysis of medical evacuation data from Africa and Papua New Guinea will be presented. Case trends show the top causes for evacuation are 1) Accident and Injury 2) Cardiovascular disease 3) Gastrointestinal disease and 4) Infectious disease. Cases in high-risk countries are more likely to result in an evacuation due to the limited local medical infrastructure, thus making the risks of deployment significantly higher in these regions. Further analysis reveals the highest risk medical conditions

to manage overseas are cardiovascular in nature. This emphasises the importance of cardiovascular risk screening in overseas fitness to work exams.

Significance of Subject Matter: Overall, the data suggests that a robust occupational health program and regular pre-travel health assessments can play a key role in attempts to reduce morbidity and mortality while abroad and reducing the medical and financial risk of managing overseas personnel. Effective prophylaxis and education can assist in mitigating the health risks posed by working overseas.

Corresponding author: Mike Broady

Email: mike.broady@internationalsos.com

The crucial role undertaken by this directorate in humanitarian assistance /civil action operations during pacific partnership 2012 and the wide scope of operations able to be managed on this platform

John McHugh

During my deployments on USNS MERCY PP08 to PNG and again in PP12 to Indonesia and the Philippines, this directorate managed a large percentage of the screened cases to be operated on aboard USNS Mercy. In my second tour during PP12 of 6 weeks, my caseload included over 20 cleft palates 18 cleft lips, many in teenagers or older children. 9 large late presenting parotid tumours, jaw tumours including an Ameloblastoma requiring resection of three quarters of the mandible, late presenting thyroid goitres and other cases such as the reconstruction of facial post traumatic deformity requiring the harvesting of a rib graft. I also saw routine lipomas and other tumours and assisted other specialities as required eg for a leg amputation. All these operations were performed aboard apart from a meningocoele through the nasoethmoidal complex in a young child which we managed in a shore based facility working with host nation surgeons. We worked closely with host nation surgeons whom we also brought aboard. The department of Surgical Services, medical, dental, optometry and veterinary directorates also spent thousands of hours engaged in SMEE, subject matter exchanges, ashore, to enhance our interoperability and facilitate good relationships with host and partner nations.

Corresponding author: John McHugh

Email: oralsurgery@optusnet.com.au

The effectiveness of civilian trauma centres and systems: lessons for the ADF?

LTCOL Michael C. Reade

After clinical training in Sydney and Melbourne, a doctorate from Oxford and an MPH from the University of Pittsburgh, Michael returned to Melbourne as an Associate Professor of Intensive Care Medicine in 2007. In 2011, after 22 years in the Army Reserve including deployments to the Balkans, Timor, the Solomon Islands and Afghanistan, he was appointed to a full-time Defence position as the Professor in Military Medicine and Surgery at the University of Queensland. He is developing a research programme at the interface of civilian and military trauma, including management of acute cognitive impairment, traumatic coagulopathy, and trauma systems design.

Modern military trauma systems evolved from structures developed in response to the conditions of the First World war. External haemorrhage control at a Regimental Aid Post in the trenches (Role 1) was followed by evacuation by stretcher bearer or motor Field Ambulance to a Casualty Clearing Station (Role 2), where (in some cases) limited lifesaving surgery could be performed in order to allow survival during transport (via road or rail) to a larger Field Hospital (Role 3). Here, definitive surgery was performed, and if return to duty was not practical, evacuation to reconstructive surgery (Role 4) and rehabilitation (Role 5) in the UK followed.

A World War One soldier would recognise all these elements in the trauma systems deployed during current and recent conflicts in the Middle East. However, modern civilian trauma medicine has evolved: major trauma is treated with simultaneous resuscitation and surgery, making 'non-surgical' resuscitation obsolete. Civilian trauma patients often bypass smaller hospitals en route to major trauma centres. They are routinely pan-CT scanned immediately after admission. Specialists in trauma surgery (as opposed to general surgery) are assisted by sub-specialists in the planned, staged management of injuries, assisted by intensive care units where patients are kept alive with the full suite of organ support (including renal replacement) in-between procedures. A modern soldier might or might not find these features in current military medical systems. Current literature provides some answers to a variety of questions relevant to the optimal design of military trauma systems, but also leaves many questions unanswered. For example,

- Do trauma centres provide better outcomes for individual patients than non-trauma centres?
- WHY do trauma centres have better outcomes?

- Do trauma systems produce better population outcomes?
- Can EVERY hospital be a trauma centre?
- Is it better to go FIRST to the closest hospital, or to a major trauma centre?
- How much longer initial transport time is it worth to get to a major trauma centre?
- Can a trauma system designed for military casualties, involving rapid evacuation to higher levels of care, also provide an adequate service to civilian casualties who cannot be evacuated?

The conclusion from the evidence suggests effective triage is the key to balancing effectiveness and efficiency in a deployed medical system. Lessons from the Middle East conflicts suggest triage can be informed by decision-support rules, but is best performed by a competent clinician next to the patient who can anticipate the interventions required, the likely course with and without such interventions, and who has the seniority and authority to design a comprehensive therapeutic plan that will be carried through immediately by the receiving hospital.

Corresponding author: Michael Reade

Email: m.reade@uq.edu.au

Human Factors

Results of the Army Colour Perception Study

Adjunct Professor John Parkes, CSC; Associate Professor Peter Nasveld; Professor Peter Warfe, CSC

John Parkes is an occupational physician with a particular interest in colour vision. He has a comprehensive colour vision testing laboratory in Melbourne and sees referred workers for colour vision testing from all over Australia. He has performed reviews of colour vision standards for various national and state organisations. He is a member of the International Colour Vision Society. He spent 19 years in the Royal Australian Navy before transferring to the Royal Australian Naval Reserve in 1996. He is a Captain and is currently Regional Director, Naval Health Reserves, for Victorian and Tasmania.

This is a study commissioned by Army and carried out by the Centre for Military and Veterans Health. A number of critical employment categories in the Australian Army were examined to determine the appropriateness of the colour perception categories that had been allotted to these categories. For many of these employment categories driving was either a primary or secondary duty. Driving in the Australian Defence Force had been considered to go beyond the standard laid down in the National Transport Commission's Assessing Fitness to Drive for Commercial and Private Vehicle Drivers 2012 (and previously 2003) and maintained the former proscription against commercial drivers with protanomalous and protanope defects promulgated in earlier National standards. The safety critical and operational critical colour vision tasks were identified for the primary and secondary duties for these employment categories, including driving tasks (including night driving, amphibious driving and operational driving). This was done through interviewing category sponsors and other experienced personnel, particularly Warrant Officers. Risk

assessments were then carried out regarding the appropriateness of the assigned colour perception categories vis-à-vis these safety critical and operational critical colour vision tasks. Out of this process recommendations were made regarding the resetting of colour perception categories as appropriate. To validate the process, two employment categories (one including driving tasks) were evaluated by observing experienced personnel from these categories both with normal colour vision and with a range of colour vision defects (that is, personnel with different colour perception categories) carrying out these safety critical and operational critical colour vision tasks. (Comprehensive colour vision testing was done on these personnel to ascertain their colour perception standards independent of the initial testing done in the course of their recruitment to the Army.) The process was thus validated.

Corresponding author: John Parkes

Email: johnparkes@chromacy.com

Exposure of aircrew to hand-held laser pointers: what are the risks?

Dr Adrian Smith, Mr David Philips

Adrian Smith is a Specialist in Aerospace Medicine and research medical officer with the RAAF Institute of Aviation Medicine and HQ Forces Command.

David Philips is a final year medical student at Australian National University who completed this project under the supervision of Dr Smith.

Exposure of aircrew to hand-held laser pointers is an increasing trend, for ADF aircrew as well as for civil aircrew around the world. Aircrew who have been exposed to lasers often experience extreme anxiety related to the perceived risks to their eyesight;

however, a review of laser events from around the world indicates that the risk is so small as to be negligible. This paper will present recent data related to exposure of aDF aircrew to lasers, and compare this to larger datasets from the commercial aviation world in order to illustrate the potential dangers of laser exposure of aircrew. Arising from the data collected, this presentation will also discuss an evidence-based approach to the management of aircrew who have been lased, in order to ensure prompt investigation and treatment of those who have suffered hazardous eye exposures but managing conservatively those who have had a benign exposure.

Corresponding author: Adrian Smith
Email: adrian@pegasusaeromed.com

Soldier Load Carriage: A investigation of the load conditioning practices of the Australian Regular Army

Assistant Professor Robin Orr, DR Rodney Pope, DR Venerina Johnston, Associate Professor Julia Coyle

Having served in the Australian Army as an infantry soldier, Physical Training Instructor, physiotherapist, and human performance officer, Rob has recently accepted a position at Bond University. Rob's fields of research include physical conditioning and injury prevention for military and protective services spanning from the initial trainee to the elite warrior. Currently focussing on tactical load carriage, Rob is exploring means of reducing injuries associated with load carriage tasks and improving the mobility and lethality of soldiers and tactical police. Published in newspapers, magazines and peer-reviewed journals, Rob is regularly invited to present at conferences both nationally and internationally.

Introduction: Soldiers must carry equipment and move, on foot, over various terrains for long and continuous periods. While the equipment carried is often crucial to mission success and survival, its weight may be a source of risk to the carrier. As physically conditioning soldiers to carry heavy load may present as a risk control strategy, the purpose of this study was to compare current Australian Regular Army (ARA) load carriage conditioning practices with previously established best practice standards for military load carriage conditioning.

Methods: ARA soldiers and units were selected via purposive sampling and invited to participate in the study. Soldier self-reported participation in load carriage physical training (PT) was captured via

online survey. In addition, textual PT programs were requested from training institutions and operational units. On receipt of textual PT programs, relevant data were extracted and each PT session rated from 1 to 4 based on the session's specificity and value to load carriage conditioning (in accordance with established best practice). A rating for each PT session was determined by reviewing the PT session title, dress, nature of the PT activity, and any clarifying comments to describe the lesson.

Results: Of the 303 survey respondents, 41% (n=126) reported participating in a load carriage PT session in the preceding fortnight. Marching Order was the most commonly reported form of dress (69%) with mean loads carried of 36.3 kg (SD=12.0kg) or 43% body weight (SD=14% BW). Roads (42%) and dirt or grass (39%) constituted the predominant terrains traversed with over 90% of sessions conducted on flat ground or over mild hills. Endurance Marching was the most common activity (60% of nominated activities) with the majority of reported sessions (79%) lasting no more than 2 hours.

The load carriage PT programs of Initial training institutions generally met with best practice evidence. However, only one (25%) of the corps training institutions built on, and progressed, these initial conditioning standards (up to 32 kg for 165 minutes of endurance marching). Two of the four trade training institutions conducted no load carriage specific conditioning over the duration of their course. Among the PT programs obtained from operational units, 50% (n=4) included load carriage specific PT (up to 31 kg for 120 minutes of endurance marching), the remaining operational units conducted no specific load carriage PT session over their PT program durations.

Conclusions: When viewed through the lense of the Frequency, Intensity, Time and Type of training principle, the load carriage practices of both the survey respondents and the conditioning programs in trade training institutions and operational units achieved limited success in meeting established evidence-based guidelines for load carriage conditioning. These findings suggest that a training gap exists between ARA current load carriage conditioning practices and the load carriage requirements of soldiers and selected army units during field training exercises and on operational duties.

Corresponding author: Robin Orr
Email: rorr@bond.edu.au

Recognition of hypoxia in a simulated flight environment.

Adrian Smith

Adrian Smith is a Specialist in Aerospace Medicine and research medical officer with the RAAF Institute of Aviation Medicine and HQ Forces Command.

Introduction: For many years hypoxia awareness training has been conducted in a benign training environment in which aircrew are required to identify their hypoxia signature whilst performing simple neurocognitive tests. This project aimed to determine if conducting hypoxia training in a simulated flight environment diminished the ability of aircrew to recognize their hypoxia signature.

Methods: 13 experienced aircrew were briefed to undertake a 45-min simulated sortie in a part-task trainer, during which hypoxia would be initiated without prior warning. The time for aircrew to recognize hypoxia symptoms and initiate recovery action was recorded, along with the SpO₂ at the time of recovery. The spectrum of symptoms and symptom severity was recorded immediately after recovery. As a baseline, hypoxia was repeated without the simulated flight task but with prior warning; elapsed time and symptoms were recorded. The baseline and

experimental conditions were compared to each other, and to the individual's memory of hypoxia from previous training.

Results: There was significant overlap in the symptoms of hypoxia experienced in the baseline and simulated flight conditions, although the simulated flight environment appeared to highlight neurocognitive and psychomotor effects more than the baseline condition. 60% of subjects initiated recovery 15 seconds earlier in the simulated flight environment than the baseline condition (paired t-test, $p < 0.03$). 69% of subjects recovered at the same level of hypoxia, with another 23% recovering at a slightly lower level of hypoxia (paired t-test, $p = 0.18$). By 2 minutes, 70% of aircrew had recovered in the simulated flight environment, compared to only 55% in the baseline condition (chi square, $p < 0.001$).

Conclusion: The results of this study suggest that aircrew may be better able to recognize hypoxia in a simulated flight environment than in a benign training setting, possibly because they are engaged in a familiar task environment within which they are able to detect subtle impairments in performance.

*Corresponding author: Adrian Smith
Email: adrian@pegasusaeromed.com*

Garrison Health

Defence medical officer attitude survey: the ADF medical employment classification review (MECR) system

CMDR Neil Westphalen

Neil graduated from the Adelaide University Medical School in 1985, and joined the RAN on 23 Jan 87 as a direct entrant. He served initially in CERBERUS before joining SWAN for an up-top deployment, followed by STALWART operating off SE Australia. He then spent a year at BNH and another at KUTTABUL (his first posting as a SMO) before proceeding on LWOP for general practice training in Launceston in 1991-2.

In Jan 1993 he joined SUCCESS for an up-top deployment as the sole Task Group MO (in company with two FFG's and a DE), followed by SYDNEY in the Northern Red Sea for DAMASK VII. He was SMO ALBATROSS in 1994-6 and joined PERTH for RIMPAC 96, thence PENGUIN for RAN Staff College Course 36/96, followed by a six month Diploma of Aviation Medicine at the RAF School of Aviation Medicine at Farnborough in the UK. During this time he also completed his FRACGP.

After another stint as SMO ALBATROSS in 1997-99, he was posted to the then HQAST as SO1 Health Ops for 11 months, then to STIRLING as SMO (and Senior Health Officer for the WA Area Health Service) from Jan 2001. Whilst SMO STIRLING he was seconded to HQ PKF in East Timor filling a RAAF AME coordinator billet; to MELVILLE for Op RELEX, and to Op GEMSBOK (to extradite VIARSA-1 from South Africa to Australia for Patagonian toothfish poaching). He also commenced his occupational medicine studies, which included a Master of Public Health.

He had a part-time posting as OIC SUMU-West and returned to the SMO STIRLING billet in Nov 2005 and completed his second fellowship a year later, as a specialist occupational and environmental physician. He was seconded to MANOORA as OIC PCRF for Ex TALISMAN SABRE in Jun 2007, and again in Nov 2008 for the SYDNEY II commemoration off Geraldton. He moved to Canberra as the Director Navy Occupational and Environmental Health in Jan 09, and took over as DNH in Sep 10. He has been the SO1 MECARS since Jun 11.

Background: There is considerable variation in the quality of the CMECR clinical documentation

received at the Medical Employment Classification Advisory and Review Services (MECARS, previously Medical Services) at Joint Health Command. This has implications regarding UMECRs and short-term fitness-for-duty decision making within the garrison health setting, noting that - unlike CMECRs - they are not subject to oversight by MECARS.

A review of civilian GP attitudes towards sickness certification in the UK and Scandinavia identified three themes, including conflict between GPs and other certification stakeholders, difficulty reconciling their various certification roles, and barriers to good practice such as their certification decision making competence. It also found that a large number of GPs would prefer not to participate in the sickness certification process.

It therefore seems likely that MECR decision making quality is significantly influenced by the attitude of the garrison health MOs who participate in the process.

Aim: This paper reports on the attitude of Defence garrison health MOs, regarding their participation in the MEC process.

Methods: A questionnaire was developed and distributed to all participants at the 20th AMMA Conference held in Melbourne in October 2011. Additional questionnaires were distributed via email through the garrison health organisation. Recipients were requested to complete a hard copy of the survey and fax it to MECARS. The results were analysed using SPSS Statistics Student Version 18.

Results: MECARS received 82 useable questionnaires from 520 garrison health MOs (15.8% response rate). Findings of note include:

- 18.3% of responders had not undertaken any MECR training, while only 29.3% of those who had undertaken the training considered it either good or very good.
- Only 54.9% rated the Member's Health Statement (MHS) and 58.5% rated the Workplace Disability Report (WDR), as either important or very important with respect to making MECR recommendations.
- Although 83.1% reported keeping copies of previous MECRs, only 22% reported using tips and tricks to reduce the time required to write MECR clinical summaries.
- Only 26.6% considered the support provided by MECARS was either good or very good.

Conclusions: The low response rate itself suggests that the overall attitude of garrison MOs towards the MECR process may not be particularly positive. However, it was still possible to draw some valid

conclusions, based on the premise that the responders most likely represent a 'best case' with respect to garrison health MO attitudes to the MECR process.

There is a need to significantly improve the quality and reach of MECR training.

There is considerable wasted effort expended on duplicating MECR information that is already available on previous MECRs.

Followup regarding the perceived MECARS support indicated frustration with the MECR process, rather than MECARS per se.

Responder's comments generally supported these conclusions. However, they also indicated that their actual level of understanding of their MECR role in practice is less than they perceive, while highlighting the need for sufficient time to conduct MECRs properly, and better IT support.

*Corresponding author: Neil Westphalen
Email: neil.westphalen1@defence.gov.au*

Nine months on: Expectation management, efficiencies and challenges for a frontline medical facility during Garrison Health Transition

SQNLDR Michael Clements

SQNLDR Clements is the Health Centre Manager at Townsville Health Centre at RAAF Base Townsville. Since completing the Graduate Medical Scheme at University of Sydney with initial training at Royal Prince Alfred Hospital in Sydney and in Geraldton, WA, he was posted to RAAF Base Tindal and then to the United Kingdom for the Diploma in Aviation Medicine. After a short period as the Officer in Charge of the 1 EHS Detachment at Townsville he was posted to his current position in Joint Health Command. Current research interests include Occupational medicine and Medical administration.

In recent years the patient dependency at RAAF Base Townsville has been serviced by two separate medical facilities, one managed by the RAAF and one by the Army. Joint Health Command assumed control of the newly named Townsville Health Centre and merged the two facilities in January 2012 and early on there were several key efficiency gains. Administrative functions, airfield emergency response liability and the cross leveling of clinical and administrative staff allowed some of the peaks and troughs experienced by the individual facilities to be balanced.

The JHC transition is a substantial game changer across the board that provides Health Centre Managers further opportunities to shape and design

the operations to suit the dependency. The experience at this facility revealed a combination of optimism and enthusiasm amongst the staff in designing the combined facility to face the new challenges and this enthusiasm has not yet waned. A key factor required to implement successful change has been the active engagement of the key stakeholders in understanding the changes and letting their own processes adapt.

Some of the ongoing concerns that remain include responding to the variable reliability of augmentees and maintaining a sense of vision, direction and leadership in the health centre staff who are heterogeneous group of contractors, APS staff, transient augmentees, non-augmentee 'supplementary' uniformed staff and JHC posted uniformed staff. Each of these staff groups have their own needs, expectations, desires and agenda's that need to be harmonised to meet Joint Health Command outcomes. With an attention to change management principles and adequate engagement of the stakeholders the restructure has provided an exciting opportunity to shape and develop an efficient, responsive and adaptable Garrison Health Service.

Corresponding author: Michael Clements

Email: michael.clements@defence.gov.au

GP Training For ADF Registrars – It Can Be Done, And It Can Be Fun

Dr Rosa Canalese, Ms Felicity Gemmell-Smith

Dr Rosa Canalese - Director of Training, GP Synergy, MBBS, Dip Paed, FRACGP, MPH, Grad Cert University Teaching

Rosa has over eighteen years of experience and a long standing interest in medical education. Whilst she has recently been involved with undergraduate medical education heading the Medical Education Unit as Associate Dean for Teaching and Learning at The University of Notre Dame, Rosa has extensive experience with vocational general practice education and training in former senior medical educator roles with the RACGP training program and the Institute of General Practice Education. She has also been a Clinical Tutor at the Universities of New South Wales and Sydney, as well as involved in undergraduate and graduate entry curriculum development. Rosa holds a Masters of Public Health and completed her Graduate Certificate in University education and teaching in 2008.

Ms Felicity Gemmell-Smith - Special Education Manager and Process Development Officer

Felicity has been involved in regional GP training since 2003, and prior to that, in nursing, administration

and community development. As part of her role with the former NEATS, she established and managed the Rural NSW GP Procedural Training Program in the New England region of NSW, which she continues to do as part of GP Synergy. In 2010 she was appointed as portfolio manager for GP Synergy's ADF registrars. She has a strong commitment to quality improvement and extensive experience in multi-disciplinary case management and project management, all of which have been extremely useful for her current roles with GP Synergy.

ADF medical personnel are able and strongly encouraged to access General Practice (GP) training through the Australian General Practice Training (AGPT) Program attached to one of 17 accredited Regional Training Providers (RTPs). To complete GP training, ADF registrars still have to fulfil all the requirements of civilian GP training whilst serving their country as full time members of the Australian Defence Force. Marrying such requirements with the conflicting demands of military vs civilian duties can cause frustration and conflict for trainees which can result in delayed training time and withdrawals from the AGPT program.

We at GP Synergy set about making GP Training easier for our ADF Registrars whilst still ensuring that they receive a comprehensive quality education program that fulfils the requirements for Fellowship of the Royal Australian College of General Practitioners (FRACGP). Through our efforts we have developed a unique training model for the delivery of the GP training components aimed at our ADF GP registrars completing their training in an average of less than 3.6 years.

GP Synergy has developed an innovative multi-disciplinary team based management approach for the ADF GP training which allows us to cope with the associated complex educational and administrative issues whilst still ensuring that our registrars continue to advance through their training despite having multiple deployments as required by their primary employer, the ADF.

Strategies we have developed to achieve this include:

- Nominated Medical Educator and Portfolio Manager for all ADF registrars within our RTP
- Demographically identified civilian practices best suited to the personality type and particular educational needs of ADF registrars that are lacking in their primary places of practice, eg geriatrics, paediatrics.
- Identified Clinical Teaching Visitors who are experienced with ADF registrars
- Set protocols and processes which allow our ADF

registrars to have deployed time counted towards their FRACGP training.

- Development of auditing and tracking tools to allow us to accurately plot where these doctors are in relation their training goals.

Achieving satisfactory exposure to the work of every GP can be a challenge for this group, and doing it in a timely manner is particularly difficult, however, we

believe that our approach is achieving adequate civilian clinical exposure whilst allowing these registrars to still service their primary employer and complete their GP training in a quality driven timely manner.

Corresponding author: Felicity Gemmill-Smith

Email: felicitygs@gpsynergy.com.au

Mental Health

Relationship between ADF member's health, partner's health, and child health outcomes: Findings from the Timor-Leste Family Study on Australian families

Anderson, R., McGuire, A., Runge, C., Breadhauer, K., Waller, M., Kanesarajah, J., Nasveld, P., & Dobson, A.

Dr Renée Anderson joined the Centre for *Military and Veterans' Health (CMVH)* in January 2011 as a Research Fellow. She has a PhD in Clinical Psychology and is particularly interested in mental health and family issues within the Australian Defence and veteran community. Other interests include PTSD, Allostatic Load, and mild traumatic brain injury for ADF members and veterans. Her clinical practice in psychology involves children, adolescents, and families.

Partners of military personnel may have elevated rates of psychiatric illness and may experience adverse physical health^[1,2]. Post-traumatic stress disorder and trauma symptoms in serving members are negatively related to marital functioning and are associated with lower relationship satisfaction for both the serving member and their partner. Further, the literature suggests that child health is correlated with the health of parents; however, studies have found the most significant predictor of child psychosocial functioning is the health of the at-home parent^[3]. In the context of Australian military deployment experiences (particularly to Timor-Leste), the current research explores the relationship between, a) ADF member's health and their partner's health, and b) ADF member, partner, and child health outcomes.

The Department of Veterans' Affairs funded a large cross-sectional quantitative study as part of the larger Family Study Program. Validated measures were used for physical (SF-12), mental (K10, SF12, PCL-C), alcohol misuse (AUDIT), family (FACES-IV), and child (SDQ) health, as well as relationship satisfaction. The sample consisted of 842 matched-partner participants, for which data was available for

both the ADF member and their partner. There were 406 families analysed, for which data was available for the ADF member, their partner, and at least one child (N = 725 children).

Key findings from the Timor-Leste Family Study suggest that physical and psychological health of the ADF member and their partner were associated. The study also found an association between the ADF member's health and their child's health, and between the at-home parent and child's health. The conclusions and implications of these findings are addressed.

References:

1. Burton, T., Farley, D., & Rhea, A. (2009). Stress-induced somatization in spouses of deployed and nondeployed servicemen. [Comparative Study; ; Research Support, Non-U.S. Gov't]. *Journal of the American Academy of Nurse Practitioners*, 21(6), 332-339.
2. Mansfield, A. J., Kaufman, J.S., Marshall, S.W., Gaynes, B.N., Morrissey, J.P., & Engel, C.C. (2010). Deployment and the use of mental health services among U.S. Army wives. *The New England Journal of Medicine*, 362(2), 101-109.
3. Al-Turkait, F., & Ohaeri, J. (2008). Post-traumatic stress disorder among wives of Kuwaiti veterans of the first Gulf War. *Journal of Anxiety Disorders*, 22(1), 18-31.

Corresponding author: Renee Anderson

Email: renee.anderson@uq.edu.au

Reservist Deployment: Perceptions of Benefits and Costs by Families and Employers

E. James Kehoe and Geoffrey J. Orme

LTCOL Kehoe came to the University of New South Wales (UNSW) as a lecturer after completing his PhD at the University of Iowa in 1976. He has conducted a productive program of basic and applied research in classical conditioning, learning, and memory. He currently is Professor and Director of Organisational

Psychology at UNSW. He also serves as a senior research consultant to the Australian Army Psychology Corps and the ADF's Joint Health Command.

The views and concerns of the families and employers of reservists from the Australian Defence Force sent on overseas deployments have not been previously surveyed. Overseas studies have been limited in size and scope. We report on surveys of family members (n = 32) and employer representatives (n = 126) of Army reservists deployed on stability operations during the period 2006-2010. In response to open-ended questions, each group generated 60% more positive statements than negative statements about aspects of the deployments for the reservists and themselves. Family members, including parents, siblings, and partners, viewed the accomplishments and service of their reservists with considerable satisfaction. The negative aspects are largely related to separation from their loved one. Support services were not needed frequently, but were highly valued when required. For the employers, the positive aspects for both their enterprise and the individual reservist employee included increases in leadership, teamwork, skills, maturity, and confidence. The negative aspects almost entirely concerned the costs associated with the absence of a valuable employee. Both surveys revealed the importance of frequent, timely communication with the ADF, as well as with the individual reservist, throughout the deployment cycle. Both groups sought information concerning the effective management of the transition of their reservist from military service back to their civilian roles.

*Corresponding author: E.James Kehoe
Email: ejameskehoe@gmail.com*

The Perceived Impact of Military Life on Children

Dr Carol Davy, Ms Michelle Lorimer, Prof Alexander McFarlane, Prof Annette Dobson

Dr Carol Davy is a Research Fellow at the University of Adelaide. She is currently an investigator on the Middle East Area of Operations Health Studies and senior researcher responsible for managing the Middle East Area of Operations Prospective Study. Dr Davy has had over 12 years experience in using both qualitative and quantitative methods to better understand the health needs of particular populations. Prior to completing her PhD in 2009, Dr Davy worked closely with the Papua New Guinea Institute of Medical Research contributing to a variety of research programs focusing on treatment seeking behaviour and evaluation studies.

Background: Deployment is a challenging time for both the individual going as well as any family

members left behind. Children, have been shown to be particularly vulnerable during this period (Kelly, Hock et al. 2001). While there is a suggestion that some cope with these challenges (Lincoln, Swift et al. 2008), many studies have found that children of deploying personnel are deeply affected by the experience (Ryan-Wenger 2001; Huebner, Mancini et al. 2007; McFarlane 2009).

Aim: This presentation examines a preliminary analysis of pre deployment data collected for the Middle East Area of Operations (MEAO) Prospective Study, in order to identify whether participants perceive that their children have been impacted by their military career.

Method:The MEAO Prospective Study is one of three integrated Military Health Outcomes Program (MilHOP) studies. All Australian Defence Force (ADF) personnel deploying to the MEAO after June 2010, and returning to Australia by May 2012 have been invited to complete a self report questionnaire prior to deploying and then again approximately three months post after returning to Australia (n~1200). A subsample of these ADF members has also undertaken a physical test (n~400) and/or a neurocognitive assessment (n~180) at both time points.

Results: Findings suggest that the odds of perceiving a negative impact on children is 1.92 times greater for participants, who have previously deployed, compared to those who have no previous deployments. In addition, both the number of deployments, and the time spent away on deployment had an affect on the perceived negative impact of deployment on children.

Conclusion: While the primary aim of the MEAO Prospective Study is to identify changes in health outcomes between pre- and post-deployment, the pre deployment dataset is already proving to be a rich source of information on not only on the physical and mental health of deploying ADF members but also on their social health including possible impacts of deployment on significant relationships.

References:

Huebner, A. J., J. A. Mancini, et al. (2007). "Parental deployment and youth in military families: exploring uncertainty and ambiguous loss." *Family Relations* 56: 112-122.

Kelly, M. L., E. Hock, et al. (2001). "Navy mothers experiencing and not experiencing deployment: reasons for staying in or leaving the military." *Military*

Psychology 13(1): 55-71.

Lincoln, A., E. Swift, et al. (2008). "Psychological adjustment and treatment of children and families with parents deployed in military combat." *Journal of Clinical Psychology* 64(8): 984-992.

McFarlane, A. C. (2009). "Military deployment: the impact on children and family adjustment and the need for care." *Current Opinion in Psychiatry* 22: 369 - 272.

Ryan-Wenger, N. A. (2001). "Impact of the threat of war on children." *American Journal of Orthopsychiatry* 71: 236-244.

Corresponding author: Carole Davy

Email: carol.davy@adelaide.edu.au

Unit cohesion, family and social support in relation to PTSD in veterans of deployments to the Gulf War, Iraq and Afghanistan:
a systematic review

Breanna Wright, Dr Helen Kelsall, Prof Malcolm Sim, Prof David Clarke, Prof Mark Creamer

Breanna Wright has a Bachelor of Arts (Honours) in psychology from the University of Melbourne and is currently undertaking a Doctor of Philosophy in the Department of Epidemiology and Preventive Medicine at Monash University. Breanna's research is part of the current follow up health study of Australian Gulf War veterans and military comparison group who were first studied in 2000-2003.

Background: Support mechanisms, such as unit cohesion, family and social support, have been associated with a range of positive health and wellbeing outcomes, particularly in deployed troops. The importance of mental health outcomes cannot be understated and one of the most common psychological morbidities reported following deployment is Posttraumatic Stress Disorder (PTSD). However, the relationship between these support mechanisms and PTSD has not been the subject of a systematic review in the veteran community.

Methods: A computer aided search was conducted of Medline, EMBASE, PubMed, Central and PsychINFO,

supplemented by searches for authors known to publish in the area and studies identified from the reference lists of included studies. Studies were eligible if participants were deployed personnel to the Gulf War, Iraq War or Afghanistan conflict, the outcome of focus was PTSD and at least one of the predictor variables of unit cohesion, family support or social support, was included. Any study design was eligible for inclusion. Two investigators systematically and independently examined all eligible studies. Quality assessment was completed via independent structured evaluation. Data extraction was performed, additional data were requested from authors where necessary and a meta-analysis was conducted.

Results: From an initial search result of 2864, 15 of these met the criteria for inclusion. Two studies utilised the same population and were counted as one study for a total of 14 included studies. The quality of the studies was generally of a high standard with acceptable response rates and use of validated instruments. A meta-analysis of unit cohesion involving six studies found that low unit cohesion was associated with PTSD caseness, with a standardised mean difference of -1.62 (-2.80, -0.45). Similarly, a meta-analysis of social support involving six studies found that low social support was associated with PTSD caseness, with a standardised mean difference of -2.40 (-3.42, -1.38). Three of the five studies on family support reported a significant relationship between low family support and PTSD caseness. Two longitudinal studies (one for social support and one for family support) indicated that longitudinally PTSD predicted low support but support did not predict later PTSD.

Conclusion: The systematic review and meta-analysis revealed that low unit cohesion, social and family support are associated with higher reporting of PTSD symptoms in returned war veterans. The review indicated that cross-sectional studies, which formed the majority of the studies eligible for the review, may be inadequate to capture the complex nature of these relationships and more longitudinal research is required.

Corresponding author: Breanna Wright

Email: breanna.wright@monash.edu

Operational Health

Evaluation of a Reintegration Presentation for Returning Australian Army Reservists

Geoffrey J. Orme, and E. James Kehoe

LTCOL Orme is an Army Reservist undertaking action research on the effect of overseas military deployments on Australian Army Reservists. This research commenced in 2002 and initially involved a two year longitudinal study of ninety Reservists deployed to East Timor in 2002-03. This research was subsequently applied in a similar fashion, using the same research tools, to three Company-size rotations of Army Reservists deployed to the Solomon Islands during 2006 and 2007 (RAMSI). Currently he is part of the Clinical Advisory Group at 3rd Health Support Battalion (Sydney, 1 HSC, Randwick Barracks, Sydney. He has led Psychological Support Teams (PST) on deployments to Bougainville, East Timor, Solomon Islands and the Middle East Area of Operations (MEAO), including Iraq.

Reservists are now widely recognised as having additional requirements for successful reintegration after an overseas deployment to their civilian lives, outside the proximal support readily available to regulars. Previous research has identified four domains in which readjustment must occur: (1) family and relationships (e.g., return from separation), (2) civilian work (e.g., career uncertainty, underemployment), (3) provider unit (e.g., reconnection, recommitment), and (4) emotional readjustment (e.g., changes in self, affective responses). Accordingly, a reservist-specific reintegration presentation was developed for delivery during the three-four days provided for decompression immediately following arrival back in Australia. This presentation was evaluated after delivery to a contingent of reservists (N = 125) who had deployed on OP ASTUTE as part of a Timor Leste Task Group (TLTG-3) for eight months from July 2011 to March 2012. The respondents (n = 82) rated the presentation as helpful, particularly, brief narratives about the experience of other reservists in their deployments to Timor Leste and the Solomon Islands. The respondents generally recommended that reservists returning from future deployments be given the presentation. The respondents also repeatedly asked that the presentation be given to their next-of-kin.

*Corresponding author: Geoffrey Orme
Email: geoff@goapl.com.au*

Sick at Sea – RAN Medevac

CMDR Ross Mills

Ross has fellowships in General Practice and Occupational and Environmental Medicine with a background in Aviation Medicine. Most recently he has worked in primarily musculoskeletal medicine and vocational rehabilitation until joining the PNF in 2009.

He is currently posted to Canberra (as a DIDO worker) as Director Navy Occupational and Environmental Health. These duties primarily involve him representing Navy's Occupational Medicine and Occupational Hygiene interests in various forums, inputting into Navy policy and providing specialist opinions on matters as they arise. Ross supplements this with regular clinical sessions and occasional medical support afloat.

In 2010 the RAN FFG, HMAS Newcastle undertook a four month deployment around the Pacific Ocean.

This presentation describes a case study of Renal Colic presenting at the worst possible time in the worst possible (geographic) location during this deployment. It is an illustration of the practice of isolated medicine without access to sophisticated monitoring equipment or treatment facilities, or extraction. This patient was transported by FFG for 7 days through the North Pacific Ocean from Hokkaido to the Aleutians, and on by air to Anchorage.

This case illustrates the difficulties faced in this scenario and emphasises the need for management to consider both worst-case medical scenarios and multiple non-medical issues. These non-medical issues included operational, communication, privacy, crew management and some of the theoretical issues relating to decision making in this context.

In the process of this Medevac records were set for the further north a RAN vessel has travelled (54° 30s N) and possibly the longest RAN medevac (1800 nm plus the air leg).

*Corresponding author: Ross Mills
Email: ross.mills2@defence.gov.au*

The Forces Command Soldier Recovery System

COL Richard Mallet

COL Richard Mallet is the Command Health Officer for Headquarters Force Command.

The development of a Soldier Recovery System in Forces Command is designed to support a Commander or soldier. Additional support is often required where complex recovery needs following wounding, injury or illness dictate the need for a more focused and comprehensive approach. Evidence suggests supporting agencies do not fully understand how to enable supported commanders in assuring the health and wellbeing of soldiers as a function of command. Enhancing situational awareness for Commanders will be an important measure of effectiveness for the Soldier Recovery System.

A Soldier Recovery Centre is the tool for Commanders in the Soldier Recovery System. Soldier Recovery Centres focus the delivery of specific and effective member-centric recovery regimes where a soldier with complex needs is identified as requiring a multi-disciplinary approach to recovery. Soldier Recovery Centres work to foster open and transparent communication between all agencies in enabling command, leadership and management functions for the chain of command. A Soldier Recovery Centre

will be required to maintain a positive recovery environment where all activities are meaningful and soldiers are enabled to focus on their recovery mission. This dictates the requirement for the development and governance of individualised and/or group recovery programs that provide a structured approach to recovery. Demonstrating the efficacy of any intervention will be an important measure of effectiveness of the Soldier Recovery System.

This paper will outline the plan for the Soldier Recovery System. It will define key criteria that continue to underpin system design. These include a system that can synchronise all agencies in supporting the recovery of wounded, injured and ill soldiers in order to ensure health and wellbeing as a function of command; a system that can govern all agencies who deliver support and services to wounded, injured and ill soldiers in demonstrating how they are having a positive impact on recovery outcomes; and a system that can facilitate education and training opportunities for soldiers whose wounding, injury or illness does not allow them to be in their normal workplace in order to keep them engaged in a positive, military environment during all stages of recovery.

Corresponding author: Richard Mallet

Email: richard.mallet@defence.gov.au

Rehabilitation

Integral Leg Prosthesis. (Early results of the osseointegration group of Australia accelerated protocol).

Al Muderis M, Bosley B, Kumar A, Laux S.

Dr. Al Muderis is an orthopaedic surgeon and a clinical lecturer at Macquarie University and The Australian School Of Advanced Medicine, specialising in Hip, Knee and Trauma surgery for the Paediatric and Adult population.

He is also the Chairman of Osseointegration Group of Australia, providing those with above-the-knee amputations with a leg replacement using the integral leg prosthesis, which is designed to be as close to the human anatomy as possible.

He is a Fellow of the Australian Orthopaedic Association and the Royal Australasian College of Surgeons.

Introduction: Transcutaneous osseointegration is an innovative technology that has been successfully used for above knee amputees since the 1990s to overcome the problems that are associated with the

standard socket prosthesis. Between 2009–2011, we operated on ten patients using this technology and we are presenting our early results.

Methods: Between 2009–2011 we performed ten procedures. This involves the insertion of a transcutaneous femoral intramedullary implant, whose most distal external aspect serves as a hard point for prosthesis attachment. There were six males and four females. Age range 23 – 58. All patients had transfemoral amputation, seven due to trauma and three due to infected total knee replacement. Preoperative assessments include medical, psychological and radiological examinations. Eight patients underwent standard two-stage procedure with a Six-week interval. Two patients had corrective surgeries prior the standard procedures. All patients were allowed early mobilization and full weight bearing two weeks after the second stage.

Results: Having studied published results of implant osseointegration in cementless total hip replacement as well as transcutaneous femoral intramedullary implant, we established the Osseointegration Group of Australia Accelerated Protocol.

Overall, there was a high level of patient satisfaction (90%). All patients except one returned to pre-amputation activities. Gait improved in all patients. All patients have retained the implant up to date. No infections to date. All patients regained osseoperception and reduced phantom pain. Skin irritation due to the old socket prosthesis has completely recovered in all patients who were using that type of prosthesis. Complications include stoma problems in one patient, one shortening of the external portion of the prosthesis due to knee joint height difference.

Discussion: The transcuteaneous femoral prosthesis is an excellent alternative and potentially will be the first choice for many transfemoral amputees in the near future. We have demonstrated that early mobilization with our accelerated protocol enables patients to regain much of their freedom in mobility without compromising the mechanical stability of the osseointegration.

References:

Aschoff HH, Kennon RE, Keggi JM, Rubin LE. Transcutaneous, distal femoral, intramedullary attachment for above-the-knee prostheses: an endo-exo device. *J Bone Joint Surg Am.* 2010 Dec;92 Suppl 2:180-6.

Branemark R. Branemark, PI. Rydevik B. Myers R. Osseointegration in skeletal reconstructions and rehabilitation: a review. *J. Rehab. Res. Dev.* 2001;(38) N2:175-181.

Corresponding author: Murjed Al Muderis

Email: murjed@me.com

Initiatives to promote successful rehabilitation outcomes and veteran wellness

Mr Simon Graham

Simon is currently a policy and research officer seeking to improve DVA's rehabilitation program and is involved in developing a range of rehabilitation policy initiatives associated with the implementation of the Military Rehabilitation and Compensation Act 2004. These have included whole of person rehabilitation reporting documentation for use by DVA providers; a comprehensive biopsychosocial approach to rehabilitation needs assessment; trialing the use of a universal assessment project and assisting development of the 'Getting the Edge' project for early intervention of complex clients in DVA. Simon has managed DVA rehabilitation research projects with

research bodies including the University of Sydney and the Australian Centre for Posttraumatic Mental Health.

The Military Rehabilitation and Compensation Act 2004 (MRCA) provides rehabilitation, treatment and compensation for injury disease and death from ADF service on or after 1 July 2004. MRCA replaces the Veterans' Entitlements Act 1986 and the Safety, Rehabilitation and Compensation Act 1988 although in some circumstances claims can still be made under these acts.

The MRCA covers permanent and reserve Australian Defence Force members, cadets, cadet officers, instructors and discharged members. The legislation mandates both the Department of Defence (for full time serving members) and the Department of Veterans' Affairs (current serving part time and discharged members) to play an integral role in the provision of rehabilitation services.

The MRCA focuses on the provision of rehabilitation services to assist members who suffer from a service-related injury or disease make as full a recovery as possible and return to their normal service duties or, after discharge, civilian work if they are able, and usual daily, family and community activities. MRCA rehabilitation provisions are consistent with the World Health Organisation's generic biopsychosocial model in management of people with injuries and provides a client centred platform from which the ADF, DVA and the service providers operate.

The Department of Veterans' Affairs delivers a wide range of rehabilitation activities for all client groups. Rehabilitation services incorporate medical, psychosocial and vocational rehabilitation activities based on the persons assessed needs.

This presentation will identify and endorse DVA's whole of department approach to rehabilitation, illustrate a number of current initiatives designed to enhance the rehabilitation experience and promote positive outcomes for the person. These include an assessment process that provides greater access to rehabilitation, the measurement of individual goals achieved and the early identification of clients potentially at risk of achieving rehabilitation goals.

Information around the On Base Advisory Service (OBAS) and how that facility can promote wellness and links with DVA's rehabilitation services will also be provided to the conference.

Corresponding author: Simon Graham

Email: simon.graham@dva.gov.au

Military History

The Magnificent Men Return!

Peter Hurly

WGDCR Peter Hurly is currently the Director of Air Force Medicine for the RNZAF. He has been a member of St John Ambulance since 1964 and was involved in ambulance work and training. He trained as a pharmacist in South Africa and studied medicine obtaining his MBChB in 1983. He worked in hospital medicine and Accident and Emergency. He was a member of the South African Military Medical Service and saw active service in South Africa. On moving to New Zealand, he joined the Royal New Zealand Army Medical Corps. He then moved to general practice in Palmerston North and took up a Reservist position with the RNZAF. He obtained a Diploma in Aviation Medicine and a certificate in Air Retrieval medicine, subsequently progressing to a Masters in Aviation medicine. He moved back into full time military medicine in 2002 and became the Director of Air Force Medicine in 2004. He is due to retire from the Regular Force at the end of this year.

Presentation surrounding the planning, logistics and support for 30 RNZAF veterans of Bomber Command, who returned to London for the unveiling of the Memorial to WW2 Bomber Command. All veterans were in their nineties and flew by RNZAF military aircraft to London and back to New Zealand for the commemoration. Back ground is given and description of the selection process, itinerary planning, personnel support and, particularly relating to the medical planning, care and support provided during the mission.

Corresponding author: PeterHurly

Email: peter.hurly@nzdf.mil.nz

A woman at war: The life and times of Dr Phoebe Chapple MM, an Australian surgeon on the Western Front.

A/Prof Susan Neuhaus, MS Sharon Mascall-Dare

Susan Neuhaus CSC served with the RAAMC in both regular and reserve roles between 1987 and 2009. She has held a variety of clinical and command appointments and is a graduate of Command and Staff College (Res). Her operational deployments include Cambodia, Bougainville and Afghanistan. Susan works in fulltime surgical practice and is actively involved in Veteran's health issues, a member of the Veterans Health Advisory Council (SA) and Chair of the Repat Foundation. Awarded a CSC in 2009, Susan was a finalist for Australian of the Year 2012.

At the outbreak of World War I female doctors were not universally accepted. Both the Australian Army and initially the British War Office, saw no role for female medical officers and refused to entertain the idea of medical woman serving in military hospitals.

Quite undeterred however, and determined to prove their medical skills, 14 of the 129 female doctors in Australia at the time found their way to European theatres of war. These women served with the Royal Army Medical Corps (RAMC) and with a variety of "All Women Medical Units". They served in France, Belgium, Greece and the Balkans, including as military surgeons. The conditions of their service and their prospects of recognition were however, quite different to those of their male counterparts.

One of these remarkable female doctors was Dr Phoebe Chapple whose experiences reflect the social and historical circumstances of the time.

Born in Adelaide on March 31 1879, she entered the University of Adelaide aged just 16, graduating with bachelor's degrees in science, surgery and medicine. Frustrated with the Australian army's refusal to appoint women doctors, she travelled at her own expense to England in February 1917 to enlist in the RAMC where she was appointed as surgeon to Cambridge Military Hospital in Aldershot. Later, she was attached to Queen Mary's Army Auxiliary Corps (QMAAC) and sent to France. Chapple was accorded the honorary rank of Captain and was one of the first two women doctors sent to the Western Front, which she 'regarded as an honor [sic] for Australia.

On 29 May 1918, Chapple was inspecting a Camp near Abbeville in France when it came under a German 'aerial bombing' attack. Her actions that night tending the wounded were recognised with the award of the Military Medal (MM). Chapple was the first woman doctor and the first Australian woman to receive such an honour "For gallantry and devotion to duty during an enemy [action]".

Chapple (now with the rank of Major) went on to serve as a doctor with the Women's Auxiliary Army Corps in Rouen and Le Havre. She returned to Adelaide in 1919 and resumed clinical practice. She died on March 24, 1967 and was cremated with full military honours.

This paper will discuss the wartime contribution of Dr Pheobe Chapple MM within the context of Australian society during WWI. Despite significant foreign awards from the governments of Britain, France, Serbia and Greece, the service of these women passed largely unnoticed within their own

country. The reasons why the wartime service of Dr Chapple and the other women doctors serving in Europe and on the Western Front has been overlooked in the official history of WWI will be explored.

*Corresponding author: Susan Neuhaus
Email: susanneuhaus@apsa.com.au*

Courage, Endurance and Care: Australian Army Mobile Hospital teams in the Kokoda Campaign

Dr Barry E Reed

Maxillofacial Surgeon John Hunter Hospital, Newcastle since 1991; Clinical Lecturer School Of Medicine, University of Newcastle; 3rd HSB; Colonel Kenny Award as best Army Reserve Dental Officer 2008 for achievements at AACAP and official visit Brooke Army Medical Center Texas regarding IED facial injuries; Award Australian Army History Research grant regarding the Kokoda Campaign; Maxillofacial Surgeon: Exercise Talisman Sabre 2009, 2011; AACAP 2011; Lecturer maxillofacial battle trauma for annual triservice course HMAS Cerberus since 2007; invited lecturer international meetings and author journal articles on maxillofacial ballistic trauma; Foundation Clinical Director Oral and Maxillofacial Surgery Unit John Hunter Hospital 1992 – 1995.

Many of the modern principles of combat health support were displayed in the critical 1942 Kokoda campaign of the Australian Army Field Ambulances and regimental aid post teams. The photographic collection of the Australian War Memorial provides a unique view of the essential and difficult work of these Field Ambulances and RAP teams and illustrate this presentation. The Field Ambulances functioned in similar fashion to a current day ADF role two (enhanced) deployable hospital providing initial wound surgery close to the battlefield while being mobile and capable of redeploying quickly to minimise evacuation times over the mountainous jungle terrain as the campaign progressed. Events of the campaign of the Field Ambulances are described in relation to current day principles of combat health support. It appears there was a lack of foresight at the time of the many great obstacles in preparation for provision of effective medical care in a combined mountain and jungle tropical campaign. These preparation shortcomings led to the need for effective improvisations, much adaptability, flexibility and considerable ingenuity by the medical soldiers of the three principal Field Ambulances involved which provided life saving solutions to the unique casualty care challenges of the Kokoda campaign and are described in this presentation. The solutions provided by these Field Ambulance soldiers and the

Brigade Headquarters staff to the challenges posed by the Kokoda Campaign included: “leapfrog” Field Ambulance movements to enable rapid casualty care; Holding non-walking casualties long term in wards; Self evacuation of the walking wounded; Aerial supply drops of urgently needed essential medical supplies; Pioneering the use of aerial casualty evacuation and the related vital role of possession of airstrips; Field training and multitasking personnel for staff shortages in key clinical roles such as providing general anaesthesia and aerial evacuation organisation; Improvisation of hospital equipment from local materials such as saplings for operating theatre tables and splints; location of medical liaison officers at Brigade Headquarters which enabled more efficient casualty care; and very importantly, the vital role of the indigenous carriers, the Fuzzy Wuzzy Angels, in resupply and casualty evacuation. Excerpts from the wartime memoirs of members of these Field Ambulances and RAP teams describing their work, their difficulties and solutions are related in this presentation. In conclusion, many of the modern principles of combat health support were displayed in the Kokoda campaign of the Field Ambulances including: mobility, proximity, flexibility, responsiveness, simplicity, continuity of care, and economy of effort. Of most importance, the timeless military medicine qualities of courage, endurance and care were magnificently exemplified by the members of these Field Ambulances and RAP teams in their Kokoda campaign.

*Corresponding author: Barry Reed
Email: bereed@bigpond.net.au*

Two different shadows: caring for Australian and American ex-POWs after World War II

Dr Rosalind Hearder

Awarded a PhD in History at the University of Melbourne, Rosalind's study of Australian medical officers' work in Japanese captivity during the Second World War was published in 2009 by Allen and Unwin. She subsequently undertook a post-doctoral Fulbright fellowship in the USA, including teaching a university course on the history of military medicine.

Rosalind has held positions in academia and government, including two years with the Official History of Australian Peacekeeping project, focusing on medical aspects such as Gulf War Syndrome and PTSD. She currently works in the Victorian Parliament, and is also writing a book based on the diaries of a British police officer in Japanese captivity.

Thousands of Australian and American troops spent

years as prisoners of the Japanese during the Second World War. Both lost over one-third of their forces through brutal starvation, disease and physical abuse. Approximately 14,000 Australian and 16,000 American surviving POWs returned home. Though allies during the war, each country would show marked difference in how survivors were treated and how their experience is remembered.

In Australia, the Pacific War was close to home and the catastrophic story of Japanese captivity became a dominant part of the larger Australian war experience. Sir Edward 'Weary' Dunlop, one of the 106 Australian medical officers in Japanese captivity, is not only the most famous POW from the Second World War, but arguably remains one of the best-known Australians from all twentieth century

conflicts. In contrast, in the United States today, many people have no idea that Japan captured any of their troops during the Pacific War, and their stories receive comparatively little attention.

While there are many reasons for this, this paper will focus on three areas: how the two groups' experiences were shaped in the immediate post-war period; the very different Australian and American Vietnam War experiences and the impact of the rise of Post-Traumatic Stress Disorder diagnoses on commemoration of the World War II POW story; and some of the consequences of failing to learn important medical lessons from Japanese captivity.

Corresponding author: Rosalind Hearder

Email: rhearder@yahoo.com